

Risk Factors of Mild and Severe Stunting Children in Rural and Urban Areas in Indonesia

by Tri Siswati

Submission date: 31-Jan-2023 11:39AM (UTC+0700)

Submission ID: 2003093620

File name: risk_factors_stunting_Q3.pdf (690.71K)

Word count: 1216

Character count: 6147



Risk Factors of Mild and Severe Stunting Children in Rural and Urban Areas in Indonesia

*Tri Siswati¹, Joko Susilo¹, Hari Kusnanto², Lukman Waris³

1. Department of Nutrition, Politeknik Kesehatan Kementerian Kesehatan, Yogyakarta, Indonesia
2. Department of Family and Community Medicine, Faculty of Medicine, Public Health and Nursing, Universitas Gadjab Mada, Yogyakarta, Indonesia
3. Department of Public Health, Universitas Faleteban, Banten, Indonesia

*Corresponding Author: Email: trisiswati14@gmail.com

(Received 11 Jan 2020; accepted 23 Jan 2020)

Dear Editor-in-Chief

Geographically, Indonesia is very wide, and therefore the disparity of stunting is different in rural, urban, or isles (1-3). Subsequently, socio-economic factors are the main cause of malnutrition (4), while distal factors such as the environment, water access, hygiene, sanitation, and housing are contributing factors (5). The proximate factors are biological aspects such as mother and children characteristics (6).

These results showed that the prevalence of stunting and severe stunting children in rural areas was higher than in urban areas, while the HAZ-score for urban children was 1.4 higher than rural children. Also, after controlling for confounding factors, low birth weight, parental stature, and economic income were the risk factors for mild and severe stunting in both urban and rural areas. Furthermore, in rural areas Water and Sanitation Hygiene (WASH) and children's age were determinant factors of stunting children, while the number of household members was a risk factor for severe stunting children. In urban areas, severe stunting was determined by the number of children. In particular, low birth weight, short parents, and economic income were the most vital risk factors

for stunting and severe stunting of children in urban and rural areas.

This research was a cross-sectional study using the pooled data from Indonesian Basic Health Research in 2007, 2010, and 2013, covering all provinces. The samples were all children 0-59 months with complete data and -5.99 to 5.99 HAZ-scores, including 38,246 children. The risk factors were characteristics of the children such as age, gender, birth weight, vitamin A supplementation. Parent's characteristics include height, education, and economic level; and household characteristics; economic level, number of household members, number of children, water, sanitation, and dwelling in rural, urban, and isles. The univariate, bivariate, and multivariate analysis used STATA-13. The highest Odds Ratio by multinomial analysis was considered the most vital risk factor.

The best multivariable analysis model found that the strongest determinant factors for stunting and severe stunting in rural and urban areas were the same, including low birth weight, short parent, and low and middle economic level (Table 1).



Table 1: Multivariate analysis of children and household characteristics related to ² stunting and severe stunting of children in urban and rural areas

Children characteristics	6 Rural		Urban	
	Severe stunting AOR (95%CI)	Stunting AOR (95%CI)	Severe stunting AOR (95%CI)	Stunting AOR (95%CI)
Age (months)				
-0-23	1	1	-	1
-24-59	1.22(1.13-1.32)	1.73(1.59-1.87)		1.31(1.21-1.42)
Sex				
-Girls	1	-	-	-
-Boys	1.14(1.06-1.23)			
Birth weight				
-Normal	1	1	1	1
-LBW	1.31(1.11-1.55)	1.46(1.24-1.72)	1.5(1.27-1.76)	1.42(1.21-1.68)
Mother's height				
-Normal	1	1	1	1
-Short	1.44(1.33-1.55)	1.64(1.51-1.77)	1.23(1.13-1.33)	1.66(1.53-1.79)
Father's height				
-Normal	1	1	1	1
-Short	1.33(1.23-1.44)	1.35(1.24-1.46)	1.36(1.24-1.48)	1.38(1.26-1.50)
Socio-demographic variables				
Number of households				
-> 9	1.66(1.07-2.56)	1.13(0.68-1.89)	-	-
-5-9	1.11(1.02-1.19)	1.13(1.05-1.23)		
-< 5	1	1		
Number of children				
->3			1.64(1.27-2.13)	-
-2-3			0.95(0.87-1.05)	
-1			1	
Father's education				
-Low	1.01(1.01-1.22)		1.09(0.98-1.22)	-
-Middle	1.06(0.95-1.18)		1.16(1.04-1.29)	
-High	1		1	
Economic level				
-Low	1.13(1.01-1.26)	1.33(1.21-1.46)	1.44(1.29-1.60)	1.42(1.28-1.57)
-Middle	1.35(1.23-1.48)	1.25(1.11-1.39)	1.17(1.07-1.29)	1.30(1.18-1.44)
-High	1	1	1	1
Mother's occupation				
-No	1.13(0.96-1.35)			1.13(0.99-1.29)
-Yes, non-formal	1.26(1.11-1.58)			1.18(1.01-1.37)
-Yes, formal	1			1
Father's occupation				
-No	-		1.13(0.86-1.48)	0.86(0.64-1.16)
-Yes, non-formal			1.17(1.07-1.29)	1.15(1.04-1.27)
-Yes, formal			1	1
WASH				
-Bad		1.45(1.12-1.87)		1.17(1.03-1.33)
-Good	-	1		1
Isles				
-Java-Bali	1	-	1	1
-Sumatra	1.26(1.14-1.38)		1.19(1.08-1.31)	1.11(1.01-1.22)
-East Indonesia	1.11(1.01-1.23)		1.11(1.01-1.23)	1.23(1.12-1.36)

In conclusion, there were no significant differences in risk factors between stunting and severe stunting among children in urban and rural areas in Indonesia. In general, low birth weight, socio-economic level, and parent's height were the risk factors for stunting and severe stunting of children in urban and rural areas. However, the number of household members was the dominant risk factor for severe stunting in rural areas, while the number of under five children was the dominant risk factor for severe stunting in urban areas. In this research, it was reported that most stunting and severe stunting children were living in unhealthy conditions with bad water access, poor hygiene and sanitation. Thus, efforts to improve the health of malnourished children focused on increasing low birth weight, economic level, WASH, family structure and family planning. Specific interventions focused on maternal and children's health to prevent baby low birth weight, while sensitive interventions focused on WASH treatment and improving economic status.

9

Conflict of interest

The authors declare that there is no conflict of interest.

References

1. NIHRD, Indonesian Ministry of Health (2007). Basic Health Research. 2007.
2. NIHRD, Indonesian Ministry of Health (2010). Basic Health Research. 2010.
3. NIHRD, Indonesian Ministry of Health (2013). Basic Health Research. 2013.
4. UNICEF (1989). The state word of children. Conceptual framework malnutrition <https://www.unicef.org/media/84766/file/SOWC-1998.pdf>
5. Torlesse H, Cronin AA, Sebayang SK, Nandy R (2016). Determinants of stunting in Indonesian children: evidence from a cross-sectional survey indicate a prominent role for the water, sanitation and hygiene sector in stunting reduction. *BMC Public Health*, 16:669.
6. WHO (2005). Commission on Social Determinants of Health, 2005-2008. <https://www.who.int/teams/social-determinants-of-health/equity-and-health/commission-on-social-determinants-of-health>

Risk Factors of Mild and Severe Stunting Children in Rural and Urban Areas in Indonesia

ORIGINALITY REPORT

21 %
SIMILARITY INDEX

13 %
INTERNET SOURCES

21 %
PUBLICATIONS

9 %
STUDENT PAPERS

PRIMARY SOURCES

1 Submitted to University of Kentucky **3** %
Student Paper

2 Md. Ashfikur Rahman, Md. Sazedur Rahman, Shakif Mohammad Shakur, Md. Hasan Howlader et al. "Risk factors of chronic childhood malnutrition: an analysis of the Bangladesh demographic and health survey 2014 data", Journal of Public Health, 2020 **3** %
Publication

3 Noor Afif Mahmudah, Dasom Im, Jeehee Pyo, Minsu Ock. "Occurrence of patient safety incidents during cancer screening: A cross-sectional investigation of the general public", Medicine, 2022 **2** %
Publication

4 eprints.poltekkesjogja.ac.id **1** %
Internet Source

5 Nurul Izzah IBRAHIM, Mohd Sharkawi AHMAD, Mohamed S ZULFARINA, Sharifah Nurul Aqilah SAYED MOHD ZARIS et al. "Risk **1** %

Factors of Osteoporosis and Post-Discharged Supplements and Drugs Intake in Patients with Lower Body Fractures", Iranian Journal of Public Health, 2021

Publication

6

cyberleninka.org

Internet Source

1 %

7

"20th International Congress of Nutrition: Granada, Spain, September 15-20, 2013", Annals of Nutrition and Metabolism, 2013

Publication

1 %

8

Bradley A. Woodruff, James P. Wirth, Ismael Ngnie-Teta, Jean Max Beaulière, Daffe Mamady, Mohamed Ag Ayoya, Fabian Rohner. "Determinants of Stunting, Wasting, and Anemia in Guinean Preschool-Age Children: An Analysis of DHS Data From 1999, 2005, and 2012", Food and Nutrition Bulletin, 2018

Publication

1 %

9

eprints.iums.ac.ir

Internet Source

1 %

10

repository.ipb.ac.id:8080

Internet Source

1 %

11

Ali JANATI, Reza EBRAHIMOGHLI, Homayoun SADEGHI-BAZARGANI, Masoumeh GHOLIZADEH, Firooz TOOFAN, Hojatolah GHARAEI. "Impact of the Iranian Health

1 %

Sector Evolution Plan on Rehospitalization: An Analysis of 158000 Hospitalizations", Iranian Journal of Public Health, 2021

Publication

12

Chenze JIAO, Shouchao WEI, Tingting LIU, Xiao BAO, Wenrong CHEN, Zhimin LIAO, Wanjuan PENG, Wei WEI, Zhou LIU. "The Prevalence of Vascular Dementia in China: A Systematic Review and Meta-Analysis from 2009-2019", Iranian Journal of Public Health, 2021

Publication

1 %

13

Raisul Akram, Marufa Sultana, Nausad Ali, Nurnabi Sheikh, Abdur Razzaque Sarker. "Prevalence and Determinants of Stunting Among Preschool Children and Its Urban-Rural Disparities in Bangladesh", Food and Nutrition Bulletin, 2018

Publication

1 %

14

www.foodandnutritionjournal.org

Internet Source

1 %

15

Siti Hawa HARITH, Norashikin MAHMUD. "The Relationship between Norms and Risky Driving Behavior: A Systematic Review", Iranian Journal of Public Health, 2020

Publication

1 %

16

Lulu'ul Badriyah, Ahmad Syafiq. "The Association Between Sanitation, Hygiene, and Stunting in Children Under Two-Years (An

1 %

Analysis of Indonesia's Basic Health Research, 2013)", Makara Journal of Health Research, 2017

Publication

17

D. Daniel. "Contextual Determinants of General Household Hygiene Conditions in Rural Indonesia", International Journal of Environmental Research and Public Health, 2021

Publication

1 %

Exclude quotes On

Exclude matches < 5 words

Exclude bibliography On