

DOKUMEN

**FULL ARTIKEL
KORESPONDENSI
ETHICAL CLEARANCE
TURNITIN**

Prevention of Cervical Cancer among Mother in Yogyakarta, Indonesia

Nanik Setiyawati, Niken Meilani
Poltekkes Kemenkes Yogyakarta, Indonesia

Article Info

Article history:

Received Sep 28, 2018

Revised Nov 8, 2018

Accepted Nov 23, 2018

Keyword:

Attitudes

Cervical cancer

Housewife

VIA test

ABSTRACT

Cervical cancer was one of the global commitments in Sustainable Development Goals (SDGs). Cervical cancer represents 7.5% of all deaths caused by cancer in women. The method to screening cervical cancer is pap smear test and Visual Inspection of Acetic Acid (VIA) test. Indonesia is the second country in the world has the most cervical cancer cases. This study aims to determine the behavior of cervical cancer prevention on housewives. This is kuantitatif research with cross sectional design. The subjects of this study were housewives in the Kota Yogyakarta and Sleman regency amounting to 350 people. The analysis used is univariate, bivariate and multivariate analysis. The results showed that there were 45.1% of mothers' have did the implementation of VIA. There is a relationship of education ($p=0.024$), level of knowledge ($p=0.036$), and maternal attitudes with cervical cancer prevention behavior. There were no correlation between age ($p=0$), parity ($p=0.816$), and family income ($p=0.174$) with the mother's behavior in the implementation of the VIA's test. Multivariate analysis showed maternal attitude ($CI=0.335-0.788$) that affect the behavior of the VIA.

Copyright © 2018 Institute of Advanced Engineering and Science.
All rights reserved.

Corresponding Author:

Nanik Setiyawati,
Midwifery Department,
Poltekkes Kemenkes Yogyakarta,
Mangkuyudan MJ III/304 Yogyakarta, Indonesia.
Email: nanikyogya@gmail.com

1. INTRODUCTION

Cervical cancer cases in 2012 included 528 000 new cases, which have been diagnosed worldwide and 85% occur in less developed areas. Cervical cancer causes 266 000 women to die of cervical cancer every year. Cervical cancer represents 7.5% of all deaths caused by cancer in women. From those data, we know that cervical cancer is a threat for the health that should be pressed down at least until 2030 which consist of curative action to the people living with cervical cancer [1].

Indonesia is the second country in the world after China has the most cervical cancer [1]. Based on data from the Ministry of Health in 2015, on average every hour the number of Cervical Cancer sufferers increased by 2.5 people and 1.1 women died of cervical cancer. The prevalence and estimation of cervical cancer patients in 2013 in Indonesia was 0.8 ‰ with an estimated total of 98,692 cases. Riau Islands Province, North Maluku Province, and D.I.Y Province (Yogyakarta Special Region) have the highest prevalence of cervical cancer which is 1.5% with a total of 2,703 cases [2].

Prevention of cervical cancer can be done by performing an early cervical health examination (screening), because symptoms of cervical cancer are not seen until the stage is more severe [3]. High-quality screening with cytology (Papanicolaou [Pap] testing) has markedly reduced mortality from squamous cell cervical cancer, which comprises 80% to 90% of cervical cancers [4-6]. Examination of early detection of

cervical cancer with VIA is a visual examination of the cervix using vinegar, meaning seeing the cervix with the naked eye to detect abnormalities after applying acetic acid or vinegar (3-5%).

To improve the implementation of prevention and early detection of cancer in women in Indonesia, the government is optimizing the cervical cancer early detection program for the 2015-2019 period. The effort taken was an early detection movement through the method of simultaneous VIA examination in all parts of Indonesia on April 21, 2015. This movement will last for five years and it is expected that by 2019 the number of women of childbearing age who are carried out early detection reaches 50 percent [2].

Mothers' awareness and preparedness with respect to the prevention of cervical cancer in their adolescent daughters were low and inadequate. Mothers should be informed and motivated to play a role in the education of their daughters regarding cervical cancer prevention. Strategies for disseminating information regarding early cervical cancer prevention for adolescent girls are recommended by communicating with both the girls and their mothers and providing them with education regarding cervical cancer prevention [7]. This study aimed to determine the factors that influence mother in behavior of cervical cancer prevention.

2. RESEARCH METHOD

This study used analytic survey with cross-sectional design. This study conducted on housewives in the Kota Yogyakarta and Sleman who were members of the Family Empowerment and Welfare amounting to 350 people. This study was conducted in 2016. The population target in this study was housewives who were members of the Family Empowerment and Welfare to manage the confounding variables. To determine the sample size, we used the formula for sample size with calculation of proportion as per Lemeshow with minimum sample size obtained was 350 respondent. Proportional sampling was used to determine the sample chosen. Variables in this study used the theoretical framework approach of Theory of Precede-Proceed (Lawrence Green) [8]. Independent variable consist of the characteristic of respondent (age, parity, education, family income), knowledge level about VIA, and mother' attitude toward VIA. The dependent variable was cervical cancer prevention behavior in mother. This study took primary data. Instrument used in this study was questionnaire. Questionnaire used was development from questionnaire about mothers' knowledge, attitude and behavior toward VIA which have been tested the validity and reliability at Bantul district.

3. RESULTS AND DISCUSSION

This study conducted on 350 respondents was housewives were members of the Family Empowerment and Welfare. The characteristic of the respondent showed in Table 1. Table 1 shows that the major proportion of respondents were aged >35 years old (76.6%), with the highest parity was secundipara in the amount of 166 respondents (47.5%). More than half of the respondents with secondary education amounted to 191 respondents (54.6%). Family income >IDR 1,450,000,00 as many as 189 respondents (54%).

Table 1. Respondent's Characteristic

Characteristic	Amount (n=350)	Percentage (%)
Age		
≤35 years	82	23.4
>35 years	268	76.6
Parity		
Nulipara	6	1.7
Primipara	83	23.7
Secundipara	166	47.5
Multi gravida	95	27.1
Education		
Basic	91	26.0
Medium	191	54.6
High	68	19.4
Family Income		
> IDR 1,450,000,00	189	54.0
≤IDR 1,450,000,00	161	46.0

Table 2 shows that most respondent have not enough knowledge about VIA (80.6%). However, there are still 35 respondents (10%) with a lack of knowledge. The role of health workers is needed in providing insight to mothers regarding early detection of cervical cancer (VIA). Most respondent have

positive attitude toward VIA (54.6%). The behavior of the VIA inspection there are 192 respondents (54.9%) for not VIA.

Table 2. Mothers' Knowledge Level, Attitude, and Behavior toward VIA Test

Characteristic	Amount (n=350)	Percentage (%)
Knowledge level about VIA		
Good	33	9.4
Adequate	282	80.6
Poor	35	10.0
Mothers' attitude toward VIA		
Positive	191	54.6
Negative	159	45.4
Mothers' behavior toward VIA		
Do VIA	158	45.1
Not Do VIA	192	54.9

Bivariate analysis was used to analyze independent and dependent variables. The relation between independent and dependent variables is showed in Table 3. Table 3 shows that factors relating to the behavior of VIA with a p-value <0.05 was educational p=0.024, p=0.036 level of knowledge, and attitudes toward VIA inspection p=0.002. 268 respondents with age>35 years as many as 147 respondents (54.9%) did not do VIA. 166 respondents with secundipara parity as many as 90 respondents (54.2%) did not do VIA. 189 respondents with family income >Rp 1.450.000,00 as many as 110 respondents (58.2%) did not do an VIA. So it can be concluded that there is no significant relationship between parity and family income with IVA behavior (p-value 0.816 and 0.174).

Table 3. The Relationship between Mothers' Characteristic, Knowledge Level, Attitude, and Behavior toward VIA

Variable	Behavior toward VIA				Total		p-value
	Not Do VIA		Do VIA		n=350	%	
	n=192	%	n=158	%			
Age							
≤35 years	45	54.9	37	45.1	82	100	
> 35 years	147	54.9	121	45.1	268	100	
Parity							
Nulipara	3	50	3	50	6	100	0.816
Primipara	48	57.8	35	42.2	83	100	
Sekundipara	90	54.2	76	45.8	166	100	
Multipara	51	53.7	44	46.3	95	100	
Education Level							
Basic	57	62.6	34	37.4	91	100	0.024*
Medium	102	53.4	89	46.6	191	100	
High	33	48.5	35	51.5	68	100	
Family Income							
> Rp.1,450,000,00	110	58.2	79	41.8	189	100	0.174
≤Rp.1,450,000,00	82	51	79	49	161	100	
Knowledge level							
Superior	16	48.5	17	51.5	33	100	0.036*
Satisfactory	150	53.2	132	46.8	282	100	
Unsatisfactory	26	74.3	9	25.7	35	100	
Attitude toward VIA							
Positive	119	62.3	72	37.7	191	100	0.002*
Negative	73	45.9	86	54.1	159	100	

Multivariate analysis were done to independent variables that have p <0.002. The result of multivariate analysis showed in Table 4. Table 4 shows that the results of statistical tests with logistic regression found that the attitude has a p-value of 0.002. This shows that the factor that most influences the behavior toward VIA is the attitude of respondents.

Cervical cancer could be prevented by screening VIA. Previous research showed that women between the age group 31 to 40 years had more awareness about cervical cancer [9]. Respondents are older than 30 years have a possibility of having a precancerous lesion is higher than under 30 years of age so that respondents in that age will find it more important to conduct tests VIA [10]. The results of this study found

no correlation of age to the prevention behavior of IVA cervical cancer ($p=0$). The results of this study are in accordance with Gustiana D's study that there was no significant relationship between age and early detection of cervical cancer $p=0.306$. It can be associated with susceptibility to disease. Age can not be used as a benchmark for a person to prevent cervical cancer. This can be due to ignorance, no complaints, nor do to prevent cervical cancer has not been necessary [11].

Table 4. Multivariate Analysis Result

	B	Sig.	Exp(B)	95.0% C.I. for EXP(B)	
				Lower	Upper
Attitude toward VIA test	-.666	.002	.514	.335	788
Constant	.164	.303	1.178		

The results of previous studies conducted by Yuliwati also showed that there was no significant relationship between age and maternal behavior to check VIA [12]. However, different research results Eva who showed that age has a significant relationship with the knowledge that motivates behavior of women for the early detection of cervical cancer by VIA in the district of Central Bogor $p=0.001$ [13].

The results of Heni P et al's research showed that parity had a significant relationship to precancerous cervical lesions [14]. While the chi square test results in this study indicate that there is no significant relationship between parity against the IVA behavior $p=0.816$. The results of this study are inconsistent with Dinengsih research and Sitanggang stated that there is a significant relationship between parity with IVA inspection. A third parity or more are at risk of developing cervical cancer. Multiparity suspected of causing a decrease in endurance. So it is necessary to do the IVA method to detect early cervical cancer [15].

The level of education a person can support or influence a person's level of knowledge and a low level of education is always related to information and knowledge is limited, the higher one's education the higher one's own understanding and knowledge of information obtained would be even higher [16]. Based on the results of this study indicate that there is a significant relationship between education with VIA behavior $p=0.024$. This research is in line with research conducted by Jia on several women in the city of Wufeng, China who explained that education affects the implementation of early detection of cervical cancer by 0.00 ($p=0.05$) with the most characteristics at the low level of education (48.4%) who are willing to carry out early detection of cervical cancer [17]. This is also confirmed by the results of further research by Eva, explains that the education factor has a significant relationship with the level of knowledge possessed by women of childbearing age, where the group of higher education have a proportion of knowledge both larger than the group of secondary education and low [13]. The level of education is one of the factors that determine a person's knowledge and perception of the importance of a case, including the importance of early detection of cervical cancer, caused by highly educated person will be broader view and more receptive to the ideas and new ways of life. It can be concluded that highly educated people will make early detection of cervical cancer [18].

The results showed that no significant relationship between family income on behavior IVA $p=0.174$. The results are consistent with research Gustiana states that there is no relationship to the economic status of cervical cancer prevention behavior ($p=0.561$), research found that respondents with lower economic status also have a good preventive health behaviors [11]. Another study conducted by Wahyuni states that there is no economic status relationship to the behavior of early detection of cervical cancer [19]. Socio-economic conditions affecting the health status change process as it affects the thoughts or beliefs that can lead to changes in health behavior. The incidence of cervical cancer is twice as large in women who have low social classes. In contrast to the results of research Febriani, women who have high incomes likely better prevention of cervical cancer, compared to women with low incomes [17].

Increased knowledge can change people's behavior from negative to positive, in addition to the knowledge also build trust [11]. Having knowledge of a disease, the concept of a disease will be formed in the individual so that it will determine a person's health behavior [20]. Lack of related knowledge about cervical cancer can be an important factor for the high incidence of cervical cancer [9]. Based on the results of chi-square test showed that there is a relationship with the level of knowledge of the behavior of IVA $p=0.036$. The results are consistent with research conducted by Yao Jia on women in the town of Wufeng China stated there is a significant relationship between knowledge with behavioral examination early detection of cervical cancer through IVA method with $p=0.000$ [17]. In addition, research conducted by Nesrin, Turkish women gained knowledge influence the early detection of cervical cancer ($p=0.001$) [21]. Based on research conducted by Khosidah, good knowledge possessed by women of childbearing age about

cervical cancer and its examination can be a motivating factor to try to avoid cervical cancer. Various sources of information that can now be easily accessed by women of childbearing age allow knowledge of cervical cancer and the examination will be better. Lack of knowledge and awareness of the importance of examination is an inhibiting factor for screening for cervical cancer [22].

Attitude is one of the personality elements that a person has to determine his actions and behavior towards an object accompanied by positive and negative feelings. Attitudes influence the formation of interest because of the tendency in the subject to accept or reject an object that is good or not [23]. The chi square test results show that there is a significant relationship between attitudes toward behavior IVA $p=0.002$. The results of this study are in line with Febriani's research the attitude of adult women in having a significant relationship with the examination of early detection of cervical cancer $p=0.025$ [18]. TRA explains that beliefs can affect attitudes and social norms which would change the form of the desire well behaved guided or happen in an individual's behavior. This theory confirms the role of one's intention in determining whether a behavior will occur [24].

4. CONCLUSION

Respondent joined this study were mostly aged >35 years old, secundipara, medium level of knowledge, family income more than minimum income regional. There is a relationship of education, level of knowledge, and attitudes with cervical cancer prevention behavior. There is no correlation between age, parity, and family income with the mother's behavior on VIA tests. Attitudes towards VIA tests is the most affect to behavior of the VIA.

ACKNOWLEDGEMENTS

We would like to thank the women and research assistants who participated in this study. The study was funded by Health Polytechnic Ministry of Health Yogyakarta, Indonesia.

REFERENCES

- [1] World Health Organization (WHO). Care of the Preterm and Low-Birth-Weight Newborn World Prematurity Day "Let Them Thrive" 1–5, 2017.
- [2] Septarini NW, Kurniati DPY, Wiryantini IAD, Putra AWGAE, Sutarga LM. "Initiation of Reproductive Health Care Village Cadres and the Early Detection of Cervical Cancer in the Village of Bangli Bali District-Inisiasi Kader Desa Peduli Kesehatan Reproduksi (KDPKR) Serta Deteksi Dini Kanker Leher Rahim Didesa Pengotan Kabupaten Bangli Bali," *Bul Udayana Mengabdikan*. 16(September):107–15, 2017.
- [3] Ministry of Health Republik Indonesia, Cancer Situation-Situasi Penyakit Kanker, Data and Information Center-InfoDATIN, 6-9, 2015.
- [4] Nordianti ME, Wahyono B. "Determinants of Visual Acetate Acid Inspection Visits in Semarang City Health Center-Determinan Kunjungan Inspeksi Visual Asam Asetat di Puskesmas Kota Semarang" *HIGEIA*, 2(1):33–44, 2018.
- [5] Gustafsson L, Ponten J, Bergstrom R, Adami HO. "International Incidence Rates of Invasive Cervical Cancer before Cytological Screening" *International Journal Cancer*. 71:159-165, 1997.
- [6] Gustafsson L, Ponten J, Zack M, Adami HO. "International Incidence Rates of Invasive Cervical Cancer After Introduction of Cytological Screening" *Cancer Causes Control*. 8:755-763, 1997.
- [7] Gustafsson L, Ponten J, Bergstrom R, Adami HO. "International Incidence Rates of Invasive Cervical Cancer Before Cytological Screening" *International Journal Cancer*. 71:159-165, 1997.
- [8] Green LW. Health Promotion Planning an Educational And Environmental Approach. Mountain View Toronto – London: Mayfield Publishing Company; 2000.
- [9] Kim HW, Kim DH. "Awareness of Cervical Cancer Prevention among Mothers of Adolescent Daughters in Korea: Qualitative Research" *BMJ Open*. 2015;1–10.
- [10] Abudukadeer A, Azam S, Mutailipu AZ, Qun L, Guilin G, Mijiti S. "Knowledge and Attitude of Uyghur Women in Xinjiang Province of China Related to the Prevention and Early Detection of Cervical Cancer" *World J Surg Oncol*. 13(110):1–7, 2015.
- [11] Mulyati S, Suwarsa O, D A IF. "Effect of Film Media on Mother's Attitude on Early Detection of Cervical Cancer-Pengaruh Media Film terhadap Sikap Ibu pada Deteksi Dini Kanker Serviks" *Jurnal Kesehatan Masyarakat*. 2015;11(2):16–24.
- [12] Gustiana D, Dewi YI, Nurchayati S, "Factors Associated with the Behavior of Prevention of Cervical Cancer in Women of Childbearing Age-Faktor-Faktor yang Berhubungan dengan Perilaku Pencegahan Kanker Serviks Pada Wanita Usia Subur" *JOM PSIK*. 2014;1(2):1–8.
- [13] Yuliwati. "Factors Associated with the Behavior of Women in Early Detection of Cervical Cancer Uterus IVA Method in Prembun Health Center 2012-Faktor-Faktor Yang Berhubungan Dengan Perilaku Wanita Dalam Deteksi Dini Kanker

- Lehar Rahim Metode IVA Di Wilayah Puskesmas Prembun 2012” 2012; Available from: <http://repository.ui.ac.id>.
- [14] Sulistiowati E. “Knowledge of Risk Factors, Behavior and Early Detection of Cervical Cancer with IVA in Women in Bogor Tengah District, Bogor-Pengetahuan tentang Faktor Risiko, Perilaku Dan Deteksi Dini Kanker Serviks dengan Inspeksi Visual Asam Asetat (IVA) pada Wanita di Kecamatan Bogor Tengah, Bogor” *Bul Penelit Kesehat.* 42(3):192–202, 2014.
- [15] Purwaningsih H, Prajatmo H. “Risk Factors Associated with the Occurrence of Precancerous Cervical Lesions at Health Centres of Karanganyar Regency” *International Journal Community Medical and Public Health.* 3(6):1437–41, 2016.
- [16] Dinengsih S, Sitanggang E. “Analysis of Early Behavior Detection of Cervical Cancer by IVA Method-Analysis Faktor Perilaku Deteksi Dini Kanker Serviks dengan Metode IVA (Inspeksi Visual Asam Asetat)” *URECOL.* 37–45, 2018.
- [17] Parapat FT, S HS, Sc M, Saraswati LD, Epid M. “Factors Associated with Behavior of Early Detection of Cervical Cancer Visual Acetate Inspection Methods in Candirot Health Center Temanggung Regency-Faktor Faktor yang Berhubungan dengan Perilaku Deteksi Dini Kanker Leher Rahim Metode Inspeksi Visual Asam Asetat di Puskesmas Candirot Kabupaten Temanggung” *Jurnal Kesehatan Masyarakat.* 4(4):363–70, 2016.
- [18] Yao J. “Knowledge about Cervical Cancer and Barriers of Screening Program among Women in Wufeng County, A High-Incidence Region of Cervical Cancer in China” 2013.
- [19] Febriani CA. “Factors Associated with Early Detection of Cervical Cancer in Gisting Subdistrict, Tanggamus Regency, Lampung-Faktor-Faktor yang Berhubungan dengan Deteksi Dini Kanker Leher Rahim di Kecamatan Gisting Kabupaten Tanggamus Lampung” *Jurnal Kesehatan.* 2015;VII(2):228–37, 2015.
- [20] Wahyuni S. “Factors Affecting Early Detection Behavior of Cervical Cancer in Ngampel District, Kendal Regency, Central Java-Faktor-Faktor yang Mempengaruhi Perilaku Deteksi Dini Kanker Serviks di Kecamatan Ngampel Kabupaten Kendal Jawa Tengah”, 2013 Available from: <https://bem.unimus.ac.id>.
- [21] Situmorang. “Relationship between Knowledge and Attitudes with Early Detection Behavior in Patients with Cervical Cancer in RSUP Dr. Kariadi Semarang in 2015-Hubungan Pengetahuan Dan Sikap Dengan Perilaku Deteksi Dini Pada Penderita Kanker Serviks Di RSUP Dr. Kariadi Semarang Tahun 2015.” *Jurnal Kesehatan Masyarakat.* 4(1):76–82, 2016.
- [22] Reis N. “Knowledge, Behavior, and Beliefs for Cervical Cancer and Screening in Turkey.” 2012; Available from: [doi:http://dx.doi.org/10.7314/APJCP.2012.13.4.1463](http://dx.doi.org/10.7314/APJCP.2012.13.4.1463)
- [23] Khosidah A. “Factors Affecting Housewives in Conducting IVA Tests as an Effort to Early Detection of Cervical Cancer-Faktor-Faktor Yang Mempengaruhi Ibu Rumah Tangga Dalam Melakukan Tes IVA Sebagai Upaya Deteksi Dini Kanker Serviks,” *Jurnal Ilmu Kebidanan.* 2015;6(2):94–105.
- [24] Sholihah A. “Relationship between the Attitude of Prevention of Cervical Cancer and Interest in Early Detection Using IVA at Fertile Women (WUS) in Mojolaban Village Sukoharjo-Hubungan Antara Sikap Pencegahan Kanker Serviks Dengan Minat Deteksi Dini Menggunakan IVA Pada WUS Di Desa Mojolaban Sukoharjo,” *Artikel Ilmiah.* 20(1), 2015.

BUKTI KORESPONDENSI
ARTIKEL JURNAL ILMIAH NASIONAL


Judul artikel : Prevention of Cervical Cancer among Mother in Yogyakarta

Jurnal : **International Journal of Public Health Science**

Penulis : Nanik Setiyawati, Niken Meilani

No	Perihal	Tanggal
1	Bukti Penerimaan artikel	28 September 2018
2	Bukti konfirmasi review dan hasil review pertama	13 Oktober 2018
3	Bukti revisi hasil review pertama	20 Oktober 2018
4	Bukti konfirmasi review dan hasil review kedua	6 November 2018
5	Bukti konfirmasi hasil review kedua dan bukti revisi hasil review kedua	8 November 2018B
6	Bukti Konfirmasi Review ketiga	9 November 2018
7	Bukti konfirmasi hasil review ketiga dan bukti revisi hasil review ketiga	10 November 2018

1. BUKTI PENERIMAAN ARTIKEL (28 SEPTEMBER 2018)



2. BUKTI KONFIRMASI REVIEW DAN HASIL REVIEW PERTAMA (13 OKTOBER 2018)

[UPHS] Editor Decision

Lina Handayani linafkm@gmail.com [uj4a2plvcpnl85798.prod.iad2.secureserver.net](mailto:linafkm@gmail.com)
to me, Niken

The following message is being delivered on behalf of International Journal of Public Health Science (IJPHS).

Nanik Setyawati:

We have reached a decision regarding your submission to International Journal of Public Health Science (IJPHS), "Prevention of Cervical Cancer on Mother in Yogyakarta, Indonesia".

Our decision is: Accepted with Revisions Required

Dr. Lina Handayani
Universitas Ahmad Dahlan
Phone +62274379418
Fax +62274381523
linafkm@gmail.com

Reviewer G:

Does the paper contain an original contribution to the field?
Yes

Is the paper technically sound?:
Yes

Does the title of the paper accurately reflect the major focus contribution of this paper?:
Yes

Please suggest change of the title as appropriate within 10 words:
-

Is the abstract a clear description of the paper?:
:
Yes

Please suggest change of the abstract:
:

Is the paper well written (clear, concise, and well organized)?:
No

Are the equations, figures and tables in this journal style, clear, relevant, and are the captions adequate?:
Yes

Please score the paper on a scale of 0 - 10 as per the directions below:
9-10 Excellent - Outstanding
7-8 Good
5-6 Average
3-4 Poor
0-2 Very Poor
:
7

Comments to the Authors (how to improve this paper):
- Re write the paper following IJPHS Template carefully
- All references must be written in English.
- Explain how did you manage confounding variables? How about reliability and validity of questionnaire?

International Journal of Public Health Science (IJPHS)
<http://www.iaescore.com/journals/index.php/IJPHS>

IJPHS_guideforauthors...

A Title Should Be the Fewest Possible Words That Accurately Describe the Content of the Paper(Center, Bold, 16pt)

First Author*, Second Author, Third Author* (10 pt)**

* Departement of Public Health Science,Ahmad Dahlan University (9 pt)

** Departement of Nutrition Health, National Chung Cheng University(9 pt)

Article Info	ABSTRACT (10 PT)
<p>Article history: Received Jun 12, 201x Revised Aug 20, 201x Accepted Aug 26, 201x</p>	<p>A well-prepared abstract enables the reader to identify the basic content of a document quickly and accurately, to determine its relevance to their interests, and thus to decide whether to read the document in its entirety. The Abstract should be informative and completely self-explanatory, provide a clear statement of the problem, the proposed approach or solution, and point out major findings and conclusions. The Abstract should be 100 to 200 words in length. The abstract should be written in the past tense. Standard nomenclature should be used and abbreviations should be avoided. No literature should be cited. The keyword list provides the opportunity to add keywords, used by the indexing and abstracting services, in addition to those already present in the title. Judicious use of keywords may increase the ease with which interested parties can locate our article (9 pt).</p> <p style="text-align: right;"><i>Copyright © 201x Institute of Advanced Engineering and Science. All rights reserved.</i></p>
<p>Keyword: First keyword Second keyword Third keyword Fourth keyword Fifth keyword</p>	
<p>Corresponding Author: Third Author, Departement of Nutrition Health, National Chung Cheng University, 168 University Road, Minhsiung Township, Chiayi County 62102, Taiwan, ROC. Email: lsntl@ccu.edu.tw</p>	

1. INTRODUCTION (10 PT)

The main text format consists of a flat left-right columns on A4 paper (quarto). The margin text from the left and top are 2.5cm, right and bottom are 2 cm. The manuscript is written in Microsoft Word, single space, Time New Roman 10pt and maximum 12 pages, which can be downloaded at the website: <http://www.iaesjournal.com/online/index.php/IJECE>

A title of article should be the fewest possible words that accurately describe the content of the paper. Omit all waste words such as "A study of ...", "Investigations of ...", "Implementation of ...", "Observations on ...", "Effect of....", "Analysis of ...", "Design of..." etc. Indexing and abstracting services depend on the accuracy of the title, extracting from it keywords useful in cross-referencing and computer

searching. An improperly titled paper may never reach the audience for which it was intended, so be specific.

The Introduction should provide a clear background, a clear statement of the problem, the relevant literature on the subject, the proposed approach or solution, and the new value of research which it is innovation. It should be understandable to colleagues from a broad range of scientific disciplines. Organization and citation of the bibliography are made in Vancouver style in sign [1], [2] and so on. The terms in foreign languages are written italic (*italic*). The text should be divided into sections, each with a separate heading and numbered consecutively. These section/subsection headings should be typed on a separate line, e.g., **1. Introduction**[3]. Authors are suggested to present their articles in the section structure: **Introduction - the comprehensive theoretical basis and/or the Proposed Method/Algorithm - Research Method - Results and Discussion – Conclusion.**

Literature review that has been done author used in the chapter "Introduction" to explain the difference of the manuscript with other papers, that it is innovative, it are used in the chapter "Research Method" to describe the step of research and used in the chapter "Results and Discussion" to support the analysis of the results [2]. If the manuscript was written really have high originality, which proposed a new method or algorithm, the additional chapter after the "Introduction" chapter and before the "Research Method" chapter can be added to explain briefly the theory and/or the proposed method/algorithm [4].

2. RESEARCH METHOD (10 PT)

Explaining research chronological, including research design, research procedure (in the form of algorithms, Pseudocode or other), how to test and data acquisition [1]-[3]. The description of the course of research should be supported references, so the explanation can be accepted scientifically [2], [4].

Tables and Figures are presented center, as shown below and cited in the manuscript.

Table 1. The Performance of ...

Variable	Speed (rpm)	Power (kW)
x	10	8.6
y	15	12.4
z	20	15.3

REFERENCES(10 PT)

The main references are international journals and proceedings. All references should be to the most pertinent and up-to-date sources. References are written in Vancouver style. Please use a consistent format for references – see examples below (9 pt):

- [1] Brambilla, M.; , "Income distribution and cause-specific mortality," *Eur J Public Health*, vol.6, no.2, pp.114-126, April 2012.
- [2] Faloutsos, M.; Karagiannis, T.; Moon, S.; , "Failure of syngeneic bone-marrow graft without preconditioning in post-hepatitis marrow aplasia," *International Journal of Public Health* , vol.24, no.5, pp.4-5, September-October 2010.
- [3] Yao-Jen Chang; Hung-Huan Liu; Tsen-Yung Wang; , "Coffee drinking and cancer of the pancreas," *Indian Journal of Public Health*, vol.16, no.3, pp.34-40, June 2009.
- [4] Chai-Arayalert, S.; Nakata, K.; , "*BN-52021 protects guinea-pig from heart anaphylaxis*," International Conference on Public Health Science 2011, vol., no., pp.220-225, 4-5 Aug. 2011.

BIOGRAPHIES OF AUTHORS (10 PT) (This part is optional)

First author's Photo (3x4cm)	Xxxx (9 pt)
Second author's photo(3x4cm)	Xxxx (9 pt)

Thirth author's
photo(3x4cm)

Xxxx (9 pt)

3. BUKTI REVISI HASIL REVIEW PERTAMA (20 OKTOBER 2018)

Prevention of Cervical Cancer on Mother in Yogyakarta, Indonesia

Nanik Setiyawati*, Niken Meilani*

*Midwifery Department Health Polytechnic Ministry of Health in Yogyakarta

Article Info	ABSTRACT
<i>Article history:</i>	Cervical cancer was one of the global commitments in Sustainable Development Goals (SDGs). Cervical cancer represents 7.5% of all deaths caused by cancer in women. The method to screening cervical cancer are pap smear test and Visual Inspection of Acetic Acid (VIA) test. Indonesia is the second country in the world has the most cervical cancer cases. This study aims to determine the behavior of cervical cancer prevention on housewives. This is kuantitatif research with cross sectional design. The subjects of this study were housewives in the Kota Yogyakarta and Sleman regency amounting to 350 people. The analysis used is univariate, bivariate and multivariate analysis. The results showed that there were 45.1% of mothers' have did the implementation of VIA. There is a relationship of education ($p = 0.024$), level of knowledge ($p = 0.036$), and maternal attitudes with cervical cancer prevention behavior. There were no correlation between age ($p = 0$), parity ($p = 0.816$), and family income ($p = 0.174$) with the mother's behavior in the implementation of the VIA's test. Multivariate analysis showed maternal attitude (CI = 0.335-0.788) that affect the behavior of the VIA.
<i>Keyword:</i> <i>VIA test</i> <i>Attitudes</i> <i>Housewife</i>	
<i>Corresponding Author:</i> Nanik Setiyawati, Midwifery Department, Helath Polytechnic Ministry of Helath Yogyakarta, Mangkuyudan MJ III/304 Yogyakarta, Indonesia. Email: nanikyogya@gmail.com	

1. INTRODUCTION

Cervical cancer cases in 2012 included 528 000 new cases, which have been diagnosed worldwide and 85% occur in less developed areas. Cervical cancer causes 266 000 women to die of cervical cancer every year. Cervical cancer represents 7.5% of all deaths caused by cancer in women. From those data, we know that cervical cancer is a threat for the health that should be pressed down at least until 2030 which consist of currative action to the people living with cervical cancer.^[1]

Indonesia is the second country in the world after China has the most cervical cancer. ^[1] Based on data from the Ministry of Health in 2015, on average every hour the number of Cervical Cancer sufferers increased by 2.5 people

and 1.1 women died of cervical cancer. The prevalence and estimation of cervical cancer patients in 2013 in Indonesia was 0.8 ‰ with an estimated total of 98,692 cases. Riau Islands Province, North Maluku Province, and D.I.Y Province (Yogyakarta Special Region) have the highest prevalence of cervical cancer which is 1.5 ‰ with a total of 2,703 cases.^[2]

Prevention of cervical cancer can be done by performing an early cervical health examination (screening), because symptoms of cervical cancer are not seen until the stage is more severe.^[3] High-quality screening with cytology (Papanicolaou [Pap] testing) has markedly reduced mortality from squamous cell cervical cancer, which comprises 80% to 90% of cervical cancers.^[4-6] Examination of early detection of cervical cancer with VIA is a visual examination of the cervix using vinegar, meaning seeing the cervix with the naked eye to detect abnormalities after applying acetic acid or vinegar (3-5%).

To improve the implementation of prevention and early detection of cancer in women in Indonesia, the government is optimizing the cervical cancer early detection program for the 2015-2019 period. The effort taken was an early detection movement through the method of simultaneous VIA examination in all parts of Indonesia on April 21, 2015. This movement will last for 5 years and it is expected that by 2019 the number of WUS (women of childbearing age) who are carried out early detection reaches 50 percent.^[2]

On the word of Hae Won Kim and Duck Hee Kim, 2015 that The mothers' awareness and preparedness with respect to the prevention of cervical cancer in their adolescent daughters were low and inadequate. Mothers should be informed and motivated to play a role in the education of their daughters regarding cervical cancer prevention. Strategies for disseminating information regarding early cervical cancer prevention for adolescent girls are recommended by communicating with both the girls and their mothers and providing them with education regarding cervical cancer prevention.^[8] This study aims to determine the factors that influence mother in behavior of cervical cancer prevention.

2. RESEARCH METHOD

This study used analytic survey with cross-sectional design. This study conducted on housewives in the Kota Yogyakarta and Sleman who were members of the Family Empowerment and Welfare amounting to 350 people. This study was conducted in 2016.

The population target in this study was housewives who were members of the Family Empowerment and Welfare to manage the confounding variables. To determine the sample size, we used the formula for sample size with calculation of proportion as per Lemeshow with minimum sample size obtained was 350 respondent. Proportional sampling was used to determine the sample chosen.

Variables in this study used the theoretical framework approach of Theory of Precede-Proceed (Lawrence Green). Independent variable consist of the characteristic of respondent (age, parity, education, family income), knowledge level about VIA, and mother' attitude toward VIA. The dependent variable was cervical cancer prevention behavior in mother.,

This study took primary data. Instrument used in this study was questionnaire. Questionnaire used was development from questionnaire about mothers' knowledge, attitude and behavior toward VIA which have been tested the validity and reliability at Bantul district.

3. RESULTS AND ANALYSIS

This study conducted on 350 respondent was housewives were members of the Family Empowerment and Welfare. The characteristic of the respondent showed in table 1 below:

Table 1. Respondent's Characteristic

Characteristic	Amount (n=350)	Percentage (%)
Age		
≤35 years	82	23.4
>35 years	268	76.6
Parity		
Nulipara	6	1.7
Primipara	83	23.7
Secundipara	166	47.5
Multi gravida	95	27.1

Education		
Basic	91	26.0
Medium	191	54.6
High	68	19.4
Family Income		
> Rp.1.450.000,00	189	54.0
≤ Rp.1.450.000,00	161	46.0

Table 1 shows that the major proportion of respondents were aged >35 years old (76.6%), with the highest parity was secundipara in the amount of 166 respondents (47.5%). More than half of the respondents with secondary education amounted to 191 respondents (54.6%). Family income > Rp. 1.450.000,00 as many as 189 respondents (54%).

Table 2. Mothers' Knowledge Level, Attitude, and Behavior toward VIA test

Characteristic	Amount (n=350)	Percentage (%)
Knowledge level about VIA		
Good	33	9.4
Adequate	282	80.6
Poor	35	10.0
Mothers' attitude toward VIA		
Positive	191	54.6
Negative	159	45.4
Mothers' behavior toward VIA		
Do VIA	158	45.1
Not Do VIA	192	54.9

Table 2 shows that most respondent have not enough knowledge about VIA (80.6%). There are still 35 respondents (10%) with a lack of knowledge, the role of health workers is needed in providing insight to mothers regarding early detection of cervical cancer (VIA). Most respondent have positive attitude toward VIA (54.6%). The behavior of the VIA inspection there are 192 respondents (54.9%) for not VIA.

Bivariate analysis was used to analyze independent and dependent variables. The relation between independent and dependent variables are showed in table 3 below:

Table 3. The Relationship between Mothers' Characteristic, Knowledge Level, Attitude, and Behavior toward VIA

Variable	Behavior toward VIA						p-value
	Not Do VIA		Do VIA		Total		
	n=192	%	n=158	%	n=350	%	
Age							
≤ 35 years	45	54.9	37	45.1	82	100	
> 35 years	147	54.9	121	45.1	268	100	
Parity							
Nulipara	3	50	3	50	6	100	
Primipara	48	57.8	35	42.2	83	100	0.816
Sekundipara	90	54.2	76	45.8	166	100	
Multipara	51	53.7	44	46.3	95	100	
Education Level							
Basic	57	62.6	34	37.4	91	100	0.024*
Medium	102	53.4	89	46.6	191	100	
High	33	48.5	35	51.5	68	100	
Family Income							
> Rp.1.450.000,00	110	58.2	79	41.8	189	100	0.174
≤ Rp.1.450.000,00	82	51	79	49	161	100	
Knowledge level							
Superior	16	48.5	17	51.5	33	100	0.036*
Satisfactory	150	53.2	132	46.8	282	100	
Unsatisfactory	26	74.3	9	25.7	35	100	
Attitude toward VIA							
Positive	119	62.3	72	37.7	191	100	0.002*
Negative	73	45.9	86	54.1	159	100	

Table 3 shows that factors relating to the behavior of VIA with a p-value <0.05 was educational p = 0.024, p = 0.036 level of knowledge, and attitudes toward VIA inspection p = 0.002. 268 respondents with age > 35 years as many as 147 respondents (54.9%) did not do VIA. 166 respondents with secundipara parity as many as 90 respondents (54.2%) did not do VIA. 189 respondents with family income >Rp 1.450.000,00 as many as 110 respondents (58.2%) did not do an VIA. So it can be concluded that there is no significant relationship between parity and family income with IVA behavior (p-value 0.816 and 0.174).

Multivariate analysis were done to independent variables that have $p < 0.002$ altogether. The result of multivariate analysis showed in table 4 below:

Table 4: Multivariate Analysis Result

	B	Sig.	Exp(B)	95.0% C.I.for EXP(B)	
				Lower	Upper
Attitude toward VIA test	-.666	.002	.514	.335	.788
Constant	.164	.303	1.178		

Table 4 shows that the results of statistical tests with logistic regression found that the attitude has a p-value of 0.002. This shows that the factor that most influences the behavior toward VIA is the attitude of respondents. Its has a p-value of $0.002 > 0.05$.

Cervical cancer could be prevented by screening VIA. The results of abudukadeer A et al (2015) showed that women between the age group 31 to 40 years had more awareness about cervical cancer.^[9] Respondents older than 30 years have a possibility of having a precancerous lesion is higher than under 30 years of age so that respondents in that age will find it more important to conduct tests VIA.^[10] The results of this study found no correlation of age to the prevention behavior of IVA cervical cancer ($p = 0$). The results of this study are in accordance with Gustiana D's (2014) study that there was no significant relationship between age and early detection of cervical cancer $p = 0.306$. It can be associated with susceptibility to disease. Age can not be used as a benchmark for a person to prevent cervical cancer. This can be due to ignorance, no complaints, nor do to prevent cervical cancer has not been necessary.^[11]

The results of previous studies conducted by Yuliwati (2012) also showed that there was no significant relationship between age and maternal behavior to check VIA.^[12] However, different research results Sulistiowati Eva (2014) who showed that age has a significant relationship with the knowledge that motivates behavior of women for the early detection of cervical cancer by VIA in the district of Central Bogor $p = 0.001$.^[13]

The results of Heni P et al's research showed that parity had a significant relationship to precancerous cervical lesions.^[14] While the chi square test results in this study indicate that there is no significant relationship between parity against the IVA behavior $p = 0.816$. The results of this study are inconsistent with Sri Dinengsih research and Erry Sitanggang (2018) states that there is a significant relationship between parity with IVA inspection. A third parity or more at risk of developing cervical cancer. Multiparity suspected of causing a decrease in endurance. So it is necessary to do the IVA method to detect early cervical cancer.^[15]

The level of education a person can support or influence a person's level of knowledge and a low level of education is always related to information and knowledge is limited, the higher one's education the higher one's own understanding and knowledge of information obtained would be even higher.^[16] Based on the results of this study indicate that there is a significant relationship between education with VIA behavior $p = 0.024$. This research is in line with research conducted by Yao Jia (2013) on several women in the city of Wufeng, China who explained that education affects the implementation of early detection of cervical cancer by 0.00 ($p = 0.05$) with the most characteristics at the low level of education (48.4%) who are willing to carry out early detection of cervical cancer.^[17] This is also confirmed by the results of further research by Sulistiowati Eva (2014), explains that the education factor has a significant relationship with the level of knowledge possessed by women of childbearing age, where the group of higher education have a proportion of knowledge both larger than the group of secondary education and low.^[13] The level of education is one of the factors that determine a person's knowledge and perception of the importance of a case, including the importance of early detection of cervical cancer, caused by highly educated person will be broader view and more receptive to the ideas and new ways of life. It can be concluded that highly educated people will make early detection of cervical cancer.^[18]

The results showed that no significant relationship between family income on behavior IVA $p = 0.174$. The results are consistent with research Gustiana (2014) states that there is no relationship to the economic status of cervical cancer prevention behavior ($p = 0.561$), research found that respondents with lower economic status also have a good preventive health behaviors.^[11] Another study conducted by Wahyuni (2013) states that there is no economic status relationship to the behavior of early detection of cervical cancer.^[19] Socio-economic conditions affecting the health status change process as it affects the thoughts or beliefs that can lead to changes in health behavior. The incidence of cervical cancer is twice as large in women who have low social classes. In contrast to the results of research Febriani CA (2016) Women who have high incomes likely better prevention of cervical cancer, compared to women with low incomes.^[17]

Increased knowledge can change people's behavior from negative to positive, in addition to the knowledge also build trust.^[11] Having knowledge of a disease, the concept of a disease will be formed in the

individual so that it will determine a person's health behavior.^[20] Lack of related knowledge about cervical cancer can be an important factor for the high incidence of cervical cancer.⁹ Based on the results of chi-square test showed that there is a relationship with the level of knowledge of the behavior of IVA $p = 0.036$. The results are consistent with research conducted by Yao Jia (2013) on women in the town of Wufeng China stated there is a significant relationship between knowledge with behavioral examination early detection of cervical cancer through IVA method with $p = 0.000$.¹⁷ In addition, research conducted by Nesrin (2012) Turkish women gained knowledge influence the early detection of cervical cancer ($p = 0.001$).^[21] Based on research conducted by Khosidah (2015), good knowledge possessed by women of childbearing age about cervical cancer and its examination can be a motivating factor to try to avoid cervical cancer. Various sources of information that can now be easily accessed by women of childbearing age allow knowledge of cervical cancer and the examination will be better. Lack of knowledge and awareness of the importance of examination is an inhibiting factor for screening for cervical cancer.^[22]

Attitude is one of the personality elements that a person has to determine his actions and behave towards an object accompanied by positive and negative feelings. Attitudes influence the formation of interest because of the tendency in the subject to accept or reject an object that is good or not.^[23] The chi square test results show that there is a significant relationship between attitudes toward behavior IVA $p = 0.002$. The results of this study are in line with Febriani CA's research (2016) the attitude of adult women in having a significant relationship with the examination of early detection of cervical cancer $p = 0.025$.^[18] TRA explains that beliefs can affect attitudes and social norms which would change the form of the desire well behaved guided or happen in an individual's behavior. This theory confirms the role of one's intention in determining whether a behavior will occur.^[24]

4. CONCLUSION

Respondent joined this study were mostly aged > 35 years old, secundipara, medium level of knowledge, family income more than minimum income regional. There is a relationship of education, level of knowledge, and attitudes with cervical cancer prevention behavior. There is no correlation between age, parity, and family income with the mother's behavior on VIA tests. Attitude towards VIA tests is the most affect to behavior of the VIA.

ACKNOWLEDGEMENTS

We would like to thank the women and research assistants who participated in this study. The study was funded by Health Polytechnic Ministry of Health Yogyakarta, Indonesia


REFERENCES

- [1] World Health Organization (WHO). Care of the preterm and low-birth-weight newborn world prematurity day - " Let Them Thrive ." 2017;1-5.
- [2] Septarini NW, Kurniati DPY, Wiryanthini IAD, Putra AWGAE, Sutarga LM. Inisiasi kader desa peduli kesehatan reproduksi (kdpkr) serta deteksi dini kanker leher rahim di desa pengotan kabupaten bangli bali. *Bul Udayana Mengabdikan*. 2017;16(September):107-15.
- [3] R I, Kementerian Kesehatan. Situasi penyakit kanker. *InfoDATIN* (2015):6-9.
- [4] Nordianti ME, Wahyono B. Determinan kunjungan inspeksi visual asam asetat di puskesmas kota semarang. *HIGEIA J PUBLIC Heal Res Dev*. 2018;2(1):33-44.
- [5] Gustafsson L, Ponten J, Bergstrom R, Adami HO. International incidence rates of invasive cervical cancer before cytological screening. *Int J Cancer*. 1997;71: 159-165.
- [6] Gustafsson L, Ponten J, Zack M, Adami HO. International incidence rates of invasive cervical cancer after introduction of cytological screening. *Cancer Causes Control*. 1997;8:755-763.
- [7] Parkin DM, Bray F. Chapter 2: The burden of HPV-related cancers. *accine*. 2006; 24(suppl 3):S3/11-25.
- [8] Gustafsson L, Ponten J, Bergstrom R, Adami HO. International incidence rates of invasive cervical cancer before cytological screening. *Int J Cancer*. 1997;71:159-165.
- [9] Kim HW, Kim DH. Awareness of cervical cancer prevention among mothers of adolescent daughters in korea : qualitative research. *BMJ Open*. 2015;1-10.
- [10] Abudukadeer A, Azam S, Mutailipu AZ, Qun L, Guilin G, Mijiti S. Knowledge and attitude of Uyghur women in Xinjiang province of China related to the prevention and early detection of cervical cancer. *World J Surg Oncol*. 2015;13(110):1-7.
- [11] Mulyati S, Suwarsa O, D A IF. Pengaruh media film terhadap sikap ibu pada deteksi dini kanker serviks. *J Kesehat Masy*. 2015;11(2):16-24.
- [12] Gustiana D, Dewi YI, Nurchayati S, Studi P, Keperawatan I, Riau U. Faktor-faktor yang berhubungan dengan perilaku pencegahan kanker serviks pada wanita usia subur. *JOM PSIK*. 2014;1(2):1-8.
- [13] Yuliwati. Faktor-faktor yang berhubungan dengan perilaku wus dalam deteksi dini kanker leher rahim metode IVA Di

Wilayah Puskesmas Prembun 2012. 2012; Available from: <http://repository.ui.ac.id>.

- [14] Sulistiowati E. Pengetahuan tentang faktor risiko, perilaku dan deteksi dini kanker serviks dengan inspeksi visual asam asetat (IVA) pada wanita di kecamatan bogor tengah. Bogor. *Bul Penelit Kesehatan*. 2014;42(3):192–202.
- [15] Purwaningsih H, Prajatmo H. Risk factors associated with the occurrence of precancerous cervical lesions at health centres of Karanganyar regency. *Int J Community Med and Public Heal*. 2016;3(6):1437–41.
- [16] Dinengsih S, Sitanggang E. Analisis faktor perilaku deteksi dini kanker serviks dengan metode IVA (Inspeksi Visual Asam Acetat). *URECOL*. 2018;37–45.
- [17] Parapat FT, S HS, Sc M, Saraswati LD, Epid M. Faktor faktor yang berhubungan dengan perilaku deteksi dini kanker leher rahim metode inspeksi visual asam asetat di puskesmas candiroto kabupaten temanggung. *J Kesehat Masy*. 2016;4(4):363–70.
- [18] Yao J. Knowledge about cervical cancer and barriers of screening program among women in wufeng county, a high-incidence region of cervical cancer in china. 2013.
- [19] Febriani CA. Faktor-faktor yang berhubungan dengan deteksi dini kanker leher rahim di kecamatan gisting kabupaten tanggamus lampung. *J Kesehatan*. 2015;VII(2):228–37.
- [20] Wahyuni S. Faktor-faktor yang mempengaruhi perilaku deteksi dini kanker serviks di kecamatan ngampel kabupaten kendal jawa tengah. 2013. Available from: <https://bem.unimus.ac.id>.
- [21] Situmorang. Hubungan pengetahuan dan sikap dengan perilaku deteksi dini pada penderita kanker serviks di RSUP dr. Kariadi semarang Tahun 2015. *J Kesehat Masy*. 2016;4(1):76–82.
- [22] Reis N. Knowledge, behavior, and beliefs for cervical cancer and screening in turkey. 2012; Available from: doi:<http://dx.doi.org/10.7314/APJCP.2012.13.4.1463>
- [23] Khosidah A. Faktor-faktor yang mempengaruhi ibu rumah tangga dalam melakukan tes IVA sebagai upaya deteksi dini kanker serviks. *J Ilm Kebidanan*. 2015;6(2):94–105.
- [24] Sholihah A. Hubungan antara sikap pencegahan kanker serviks dengan minat deteksi dini menggunakan IVA pada WUS di desa mojolaban sukoharjo. *Artik Ilm*. 2015;20(1).
- [25] Green LW. *Health promotion planning an educational and environmental approach*. Mountain View Toronto - London: Mayfield Publishing Company; 2000.

4. BUKTI KONFIRMASI REVIEW KEDUA (6 NOVEMBER 2018)




The screenshot shows an email interface with a left sidebar and a main content area. The sidebar includes sections for 'Compose', 'Inbox' (383), 'Starred', 'Snoozed', 'Important', 'Sent', 'Drafts' (55), 'Categories', 'Junk', 'More', 'Meet', 'New meeting', 'Join a meeting', and 'Hangouts' (Nanik +). The main content area displays two email messages.

Message 1:
Subject: [JPHS] Please Revise Your Paper
From: Azidanti Saufi <ufi.laes@gmail.com>
To: me, nikenbundaqueena
Date: Tue, Nov 6, 2018, 9:07 AM
Content: Dear Authors, For preparing your final paper to publish in International Journal of Public Health Science (JPHS) issue December 2018, Please revise your last paper based on the editor's comments or reviewers on the website. You have 3 days, until November 09, 2018 to revised your paper. Please submit your revised paper by reply this email. Thank you for your cooperation.
Best Regards
Azidanti Saufi
Administrator on behalf Managing Editor
Institute of Advanced Engineering and Science (IAES)
e-mail: ufi.laes@gmail.com

Message 2:
From: Nanik Setiyawati <nanikyogya@gmail.com>
To: Azidanti, nikenbundaqueena
Date: Tue, Nov 6, 2018, 9:11 AM
Content: Thank you for your information, i will do that.
Best regards
Nanik Setiyawati

5. BUKTI KONFIRMASI HASIL REVIEW KEDUA DAN BUKTI REVISI HASIL REVIEW KEDUA (8 NOVEMBER 2018)



Prevention of Cervical Cancer on Mother in Yogyakarta, Indonesia

Nanik Setiyawati*, Niken Meilani*

*Midwifery Department Health Polytechnic Ministry of Health in Yogyakarta

Article Info	ABSTRACT
<i>Article history:</i>	Cervical cancer was one of the global commitments in Sustainable Development Goals (SDGs). Cervical cancer represents 7.5% of all deaths caused by cancer in women. The method to screening cervical cancer are pap smear test and Visual Inspection of Acetic Acid (VIA) test. Indonesia is the second country in the world has the most cervical cancer cases. This study aims to determine the behavior of cervical cancer prevention on housewives. This is kuantitatif research with cross sectional design. The subjects of this study were housewives in the Kota Yogyakarta and Sleman regency amounting to 350 people. The analysis used is univariate, bivariate and multivariate analysis. The results showed that there were 45.1% of mothers' have did the implementation of VIA. There is a relationship of education ($p = 0.024$), level of knowledge ($p = 0.036$), and maternal attitudes with cervical cancer prevention behavior. There were no correlation between age ($p = 0$), parity ($p = 0.816$), and family income ($p = 0.174$) with the mother's behavior in the implementation of the VIA's test. Multivariate analysis showed maternal attitude (CI = 0.335-0.788) that affect the behavior of the VIA.
<i>Keyword:</i>	
<i>VIA test</i>	
<i>Attitudes</i>	
<i>Housewife</i>	
<i>Corresponding Author:</i>	
Nanik Setiyawati, Midwifery Department, Helath Polytechnic Ministry of Helath Yogyakarta, Mangkuyudan MJ III/304 Yogyakarta, Indonesia. Email: nanikyogya@gmail.com	

5. INTRODUCTION

Cervical cancer cases in 2012 included 528 000 new cases, which have been diagnosed worldwide and 85% occur in less developed areas. Cervical cancer causes 266 000 women to die of cervical cancer every year. Cervical cancer represents 7.5% of all deaths caused by cancer in women. From those data, we know that cervical cancer is a threat for the health that should be pressed down at least until 2030 which consist of currative action to the people living with cervical cancer.^[1]

Indonesia is the second country in the world after China has the most cervical cancer.^[1] Based on data from the Ministry of Health in 2015, on average every hour the number of Cervical Cancer sufferers increased by 2.5 people and 1.1 women died of cervical cancer. The prevalence and estimation of cervical cancer patients in 2013 in Indonesia was 0.8 ‰ with an estimated total of 98,692 cases. Riau Islands Province, North Maluku Province, and D.I.Y Province (Yogyakarta Special Region) have the highest prevalence of cervical cancer which is 1.5 ‰ with a total of 2,703 cases.^[2]

Prevention of cervical cancer can be done by performing an early cervical health examination (screening), because symptoms of cervical cancer are not seen until the stage is more severe.^[3] High-quality screening with cytology (Papanicolaou [Pap] testing) has markedly reduced mortality from squamous cell cervical cancer, which comprises 80% to 90% of cervical cancers.^[4-6] Examination of early detection of cervical cancer with VIA is a visual

examination of the cervix using vinegar, meaning seeing the cervix with the naked eye to detect abnormalities after applying acetic acid or vinegar (3-5%).

To improve the implementation of prevention and early detection of cancer in women in Indonesia, the government is optimizing the cervical cancer early detection program for the 2015-2019 period. The effort taken was an early detection movement through the method of simultaneous VIA examination in all parts of Indonesia on April 21, 2015. This movement will last for 5 years and it is expected that by 2019 the number of WUS (women of childbearing age) who are carried out early detection reaches 50 percent.^[2]

On the word of Hae Won Kim and Duck Hee Kim, 2015 that The mothers' awareness and preparedness with respect to the prevention of cervical cancer in their adolescent daughters were low and inadequate. Mothers should be informed and motivated to play a role in the education of their daughters regarding cervical cancer prevention. Strategies for disseminating information regarding early cervical cancer prevention for adolescent girls are recommended by communicating with both the girls and their mothers and providing them with education regarding cervical cancer prevention.^[8] This study aims to determine the factors that influence mother in behavior of cervical cancer prevention.

6. RESEARCH METHOD

This study used analytic survey with cross-sectional design. This study conducted on housewives in the Kota Yogyakarta and Sleman who were members of the Family Empowerment and Welfare amounting to 350 people. This study was conducted in 2016.

The population target in this study was housewives who were members of the Family Empowerment and Welfare to manage the confounding variables. To determine the sample size, we used the formula for sample size with calculation of proportion as per Lemeshow with minimum sample size obtained was 350 respondent. Proportional sampling was used to determine the sample chosen.

Variables in this study used the theoretical framework approach of Theory of Precede-Proceed (Lawrence Green). Independent variable consist of the characteristic of respondent (age, parity, education, family income), knowledge level about VIA, and mother' attitude toward VIA. The dependent variable was cervical cancer prevention behavior in mother.,

This study took primary data. Instrument used in this study was questionnaire. Questionnaire used was development from questionnaire about mothers' knowledge, attitude and behavior toward VIA which have been tested the validity and reliability at Bantul district.

7. RESULTS AND ANALYSIS

This study conducted on 350 respondent was housewives were members of the Family Empowerment and Welfare. The characteristic of the respondent showed in table 1 below:

Table 1. Respondent's Characteristic

Characteristic	Amount (n=350)	Percentage (%)
Age		
≤35 years	82	23.4
>35 years	268	76.6
Parity		
Nulipara	6	1.7
Primipara	83	23.7
Secundipara	166	47.5
Multi gravida	95	27.1
Education		
Basic	91	26.0
Medium	191	54.6
High	68	19.4
Family Income		
> Rp.1.450.000,00	189	54.0
≤ Rp.1.450.000,00	161	46.0

Table 1 shows that the major proportion of respondents were aged >35 years old (76.6%), with the highest parity was secundipara in the amount of 166 respondents (47.5%). More than half of the respondents with secondary education amounted to 191 respondents (54.6%). Family income > Rp. 1.450.000,00 as many as 189 respondents (54%).

Table 2. Mothers' Knowledge Level, Attitude, and Behavior toward VIA test

Characteristic	Amount (n=350)	Percentage (%)
Knowledge level about VIA		
Good	33	9.4
Adequate	282	80.6
Poor	35	10.0
Mothers' attitude toward VIA		
Positive	191	54.6
Negative	159	45.4
Mothers' behavior toward VIA		
Do VIA	158	45.1
Not Do VIA	192	54.9

Table 2 shows that most respondent have not enough knowledge about VIA (80.6%). There are still 35 respondents (10%) with a lack of knowledge, the role of health workers is needed in providing insight to mothers regarding early detection of cervical cancer (VIA). Most respondent have positive attitude toward VIA (54.6%). The behavior of the VIA inspection there are 192 respondents (54.9%) for not VIA.

Bivariate analysis was used to analyze independent and dependent variables. The relation between independent and dependent variables are showed in table 3 below:

Table 3. The Relationship between Mothers' Characteristic, Knowledge Level, Attitude, and Behavior toward VIA

Variable	Behavior toward VIA						p-value
	Not Do VIA		Do VIA		Total		
	n=192	%	n=158	%	n=350	%	
Age							
≤ 35 years	45	54.9	37	45.1	82	100	
> 35 years	147	54.9	121	45.1	268	100	
Parity							
Nulipara	3	50	3	50	6	100	
Primipara	48	57.8	35	42.2	83	100	0.816
Sekundipara	90	54.2	76	45.8	166	100	
Multipara	51	53.7	44	46.3	95	100	
Education Level							
Basic	57	62.6	34	37.4	91	100	0.024*
Medium	102	53.4	89	46.6	191	100	
High	33	48.5	35	51.5	68	100	
Family Income							
> Rp.1.450.000,00	110	58.2	79	41.8	189	100	0.174
≤ Rp.1.450.000,00	82	51	79	49	161	100	
Knowledge level							
Superior	16	48.5	17	51.5	33	100	0.036*
Satisfactory	150	53.2	132	46.8	282	100	
Unsatisfactory	26	74.3	9	25.7	35	100	
Attitude toward VIA							
Positive	119	62.3	72	37.7	191	100	0.002*
Negative	73	45.9	86	54.1	159	100	

Table 3 shows that factors relating to the behavior of VIA with a p-value <0.05 was educational p = 0.024, p = 0.036 level of knowledge, and attitudes toward VIA inspection p = 0.002. 268 respondents with age > 35 years as many as 147 respondents (54.9%) did not do VIA. 166 respondents with secundipara parity as many as 90 respondents (54.2%) did not do VIA. 189 respondents with family income >Rp 1.450.000,00 as many as 110 respondents (58.2%) did not do an VIA. So it can be concluded that there is no significant relationship between parity and family income with IVA behavior (p-value 0.816 and 0.174).

Multivariate analysis were done to independent variables that have $p < 0.002$ altogether. The result of multivariate analysis showed in table 4 below:

Table 4: Multivariate Analysis Result

	B	Sig.	Exp(B)	95.0% C.I.for EXP(B)	
				Lower	Upper
Attitude toward VIA test	-.666	.002	.514	.335	.788
Constant	.164	.303	1.178		

Table 4 shows that the results of statistical tests with logistic regression found that the attitude has a p-value of 0.002. This shows that the factor that most influences the behavior toward VIA is the attitude of respondents. Its has a p-value of $0.002 > 0.05$.

Cervical cancer could be prevented by screening VIA. The results of abudukadeer A et al (2015) showed that women between the age group 31 to 40 years had more awareness about cervical cancer.^[9] Respondents older than 30 years have a possibility of having a precancerous lesion is higher than under 30 years of age so that respondents in that age will find it more important to conduct tests VIA.^[10] The results of this study found no correlation of age to the prevention behavior of IVA cervical cancer ($p = 0$). The results of this study are in accordance with Gustiana D's (2014) study that there was no significant relationship between age and early detection of cervical cancer $p = 0.306$. It can be associated with susceptibility to disease. Age can not be used as a benchmark for a person to prevent cervical cancer. This can be due to ignorance, no complaints, nor do to prevent cervical cancer has not been necessary.^[11]

The results of previous studies conducted by Yuliwati (2012) also showed that there was no significant relationship between age and maternal behavior to check VIA.^[12] However, different research results Sulistiowati Eva (2014) who showed that age has a significant relationship with the knowledge that motivates behavior of women for the early detection of cervical cancer by VIA in the district of Central Bogor $p = 0.001$.^[13]

The results of Heni P et al's research showed that parity had a significant relationship to precancerous cervical lesions.^[14] While the chi square test results in this study indicate that there is no significant relationship between parity against the IVA behavior $p = 0.816$. The results of this study are inconsistent with Sri Dinengsih research and Erry Sitanggang (2018) states that there is a significant relationship between parity with IVA inspection. A third parity or more at risk of developing cervical cancer. Multiparity suspected of causing a decrease in endurance. So it is necessary to do the IVA method to detect early cervical cancer.^[15]

The level of education a person can support or influence a person's level of knowledge and a low level of education is always related to information and knowledge is limited, the higher one's education the higher one's own understanding and knowledge of information obtained would be even higher.^[16] Based on the results of this study indicate that there is a significant relationship between education with VIA behavior $p = 0.024$. This research is in line with research conducted by Yao Jia (2013) on several women in the city of Wufeng, China who explained that education affects the implementation of early detection of cervical cancer by 0.00 ($p = 0.05$) with the most characteristics at the low level of education (48.4%) who are willing to carry out early detection of cervical cancer.^[17] This is also confirmed by the results of further research by Sulistiowati Eva (2014), explains that the education factor has a significant relationship with the level of knowledge possessed by women of childbearing age, where the group of higher education have a proportion of knowledge both larger than the group of secondary education and low.^[13] The level of education is one of the factors that determine a person's knowledge and perception of the importance of a case, including the importance of early detection of cervical cancer, caused by highly educated person will be broader view and more receptive to the ideas and new ways of life. It can be concluded that highly educated people will make early detection of cervical cancer.^[18]

The results showed that no significant relationship between family income on behavior IVA $p = 0.174$. The results are consistent with research Gustiana (2014) states that there is no relationship to the economic status of cervical cancer prevention behavior ($p = 0.561$), research found that respondents with lower economic status also have a good preventive health behaviors.^[11] Another study conducted by Wahyuni (2013) states that there is no economic status relationship to the behavior of early detection of cervical cancer.^[19] Socio-economic conditions affecting the health status change process as it affects the thoughts or beliefs that can lead to changes in health behavior. The incidence of cervical cancer is twice as large in women who have low social classes. In contrast to the results of research Febriani CA (2016) Women who have high incomes likely better prevention of cervical cancer, compared to women with low incomes.^[17]

Increased knowledge can change people's behavior from negative to positive, in addition to the knowledge also build trust.^[11] Having knowledge of a disease, the concept of a disease will be formed in the

individual so that it will determine a person's health behavior.^[20] Lack of related knowledge about cervical cancer can be an important factor for the high incidence of cervical cancer.⁹ Based on the results of chi-square test showed that there is a relationship with the level of knowledge of the behavior of IVA $p = 0.036$. The results are consistent with research conducted by Yao Jia (2013) on women in the town of Wufeng China stated there is a significant relationship between knowledge with behavioral examination early detection of cervical cancer through IVA method with $p = 0.000$.¹⁷ In addition, research conducted by Nesrin (2012) Turkish women gained knowledge influence the early detection of cervical cancer ($p = 0.001$).^[21] Based on research conducted by Khosidah (2015), good knowledge possessed by women of childbearing age about cervical cancer and its examination can be a motivating factor to try to avoid cervical cancer. Various sources of information that can now be easily accessed by women of childbearing age allow knowledge of cervical cancer and the examination will be better. Lack of knowledge and awareness of the importance of examination is an inhibiting factor for screening for cervical cancer.^[22]

Attitude is one of the personality elements that a person has to determine his actions and behave towards an object accompanied by positive and negative feelings. Attitudes influence the formation of interest because of the tendency in the subject to accept or reject an object that is good or not.^[23] The chi square test results show that there is a significant relationship between attitudes toward behavior IVA $p = 0.002$. The results of this study are in line with Febriani CA's research (2016) the attitude of adult women in having a significant relationship with the examination of early detection of cervical cancer $p = 0.025$.^[18] TRA explains that beliefs can affect attitudes and social norms which would change the form of the desire well behaved guided or happen in an individual's behavior. This theory confirms the role of one's intention in determining whether a behavior will occur.^[24]

8. CONCLUSION

Respondent joined this study were mostly aged > 35 years old, secundipara, medium level of knowledge, family income more than minimum income regional. There is a relationship of education, level of knowledge, and attitudes with cervical cancer prevention behavior. There is no correlation between age, parity, and family income with the mother's behavior on VIA tests. Attitude towards VIA tests is the most affect to behavior of the VIA.

ACKNOWLEDGEMENTS


We would like to thank the women and research assistants who participated in this study. The study was funded by Health Polytechnic Ministry of Health Yogyakarta, Indonesia

REFERENCES

- [26] World Health Organization (WHO). Care of the preterm and low-birth-weight newborn world prematurity day - " Let Them Thrive ." 2017;1-5.
- [27] Septarini NW, Kurniati DPY, Wiryanthini IAD, Putra AWGAE, Sutarga LM. Inisiasi kader desa peduli kesehatan reproduksi (kdpkr) serta deteksi dini kanker leher rahim di desa pengotan kabupaten bangli bali. *Bul Udayana Mengabdi*. 2017;16(September):107-15.
- [28] R I, Kementerian Kesehatan. Situasi penyakit kanker. *InfoDATIN* (2015):6-9.
- [29] Nordianti ME, Wahyono B. Determinan kunjungan inspeksi visual asam asetat di puskesmas kota semarang. *HIGEIA J PUBLIC Heal Res Dev*. 2018;2(1):33-44.
- [30] Gustafsson L, Ponten J, Bergstrom R, Adami HO. International incidence rates of invasive cervical cancer before cytological screening. *Int J Cancer*. 1997;71: 159-165.
- [31] Gustafsson L, Ponten J, Zack M, Adami HO. International incidence rates of invasive cervical cancer after introduction of cytological screening. *Cancer Causes Control*. 1997;8:755-763.
- [32] Parkin DM, Bray F. Chapter 2: The burden of HPV-related cancers. *accine*. 2006; 24(suppl 3):S3/11-25.
- [33] Gustafsson L, Ponten J, Bergstrom R, Adami HO. International incidence rates of invasive cervical cancer before cytological screening. *Int J Cancer*. 1997;71:159-165.
- [34] Kim HW, Kim DH. Awareness of cervical cancer prevention among mothers of adolescent daughters in Korea : qualitative research. *BMJ Open*. 2015;1-10.
- [35] Abudukadeer A, Azam S, Mutailipu AZ, Qun L, Guilin G, Mijiti S. Knowledge and attitude of Uyghur women in Xinjiang province of China related to the prevention and early detection of cervical cancer. *World J Surg Oncol*. 2015;13(110):1-7.
- [36] Mulyati S, Suwarsa O, D A IF. Pengaruh media film terhadap sikap ibu pada deteksi dini kanker serviks. *J Kesehat Masy*. 2015;11(2):16-24.
- [37] Gustiana D, Dewi YI, Nurchayati S, Studi P, Keperawatan I, Riau U. Faktor-faktor yang berhubungan dengan perilaku pencegahan kanker serviks pada wanita usia subur. *JOM PSIK*. 2014;1(2):1-8.
- [38] Yuliwati. Faktor-faktor yang berhubungan dengan perilaku wus dalam deteksi dini kanker leher rahim metode IVA Di

- Wilayah Puskesmas Prembun 2012. 2012; Available from: <http://repository.ui.ac.id>.
- [39] Sulistiowati E. Pengetahuan tentang faktor risiko, perilaku dan deteksi dini kanker serviks dengan inspeksi visual asam asetat (IVA) pada wanita di kecamatan bogor tengah. Bogor. *Bul Penelit Kesehat*. 2014;42(3):192–202.
- [40] Purwaningsih H, Prajatmo H. Risk factors associated with the occurrence of precancerous cervical lesions at health centres of Karanganyar regency. *Int J Community Med and Public Heal*. 2016;3(6):1437–41.
- [41] Dinengsih S, Sitanggang E. Analisis faktor perilaku deteksi dini kanker serviks dengan metode IVA (Inspeksi Visual Asam Acetat). *URECOL*. 2018;37–45.
- [42] Parapat FT, S HS, Sc M, Saraswati LD, Epid M. Faktor faktor yang berhubungan dengan perilaku deteksi dini kanker leher rahim metode inspeksi visual asam asetat di puskesmas candirot kabupaten temanggung. *J Kesehat Masy*. 2016;4(4):363–70.
- [43] Yao J. Knowledge about cervical cancer and barriers of screening program among women in wufeng county, a high-incidence region of cervical cancer in china. 2013.
- [44] Febriani CA. Faktor-faktor yang berhubungan dengan deteksi dini kanker leher rahim di kecamatan gisting kabupaten tanggamus lampung. *J Kesehat*. 2015;VII(2):228–37.
- [45] Wahyuni S. Faktor-faktor yang mempengaruhi perilaku deteksi dini kanker serviks di kecamatan ngampel kabupaten kendal jawa tengah. 2013. Available from: <https://bem.unimus.ac.id>.
- [46] Situmorang. Hubungan pengetahuan dan sikap dengan perilaku deteksi dini pada penderita kanker serviks di RSUP dr. Kariadi semarang Tahun 2015. *J Kesehat Masy*. 2016;4(1):76–82.
- [47] Reis N. Knowledge, behavior, and beliefs for cervical cancer and screening in turkey. 2012; Available from: doi:<http://dx.doi.org/10.7314/APJCP.2012.13.4.1463>
- [48] Khosidah A. Faktor-faktor yang mempengaruhi ibu rumah tangga dalam melakukan tes IVA sebagai upaya deteksi dini kanker serviks. *J Ilm Kebidanan*. 2015;6(2):94–105.
- [49] Sholihah A. Hubungan antara sikap pencegahan kanker serviks dengan minat deteksi dini menggunakan IVA pada WUS di desa mojolaban sukoharjo. *Artik Ilm*. 2015;20(1).
- [50] Green LW. *Health promotion planning an educational and environmental approach*. Mountain View Toronto - London: Mayfield Publishing Company; 2000.

6. BUKTI KONFIRMASI REVIEW KETIGA (9 NOVEMBER 2018)



Prevention of Cervical Cancer on Mother in Yogyakarta, Indonesia

Nanik Setiyawati, Niken Meilani

Midwifery Department Health Polytechnic Ministry of Health in Yogyakarta

Article Info

Article history:

Received Sep 28, 2018

Revised Nov 8, 2018

Accepted Nov 15, 2018

Keyword:

Attitudes
Housewife
VIA test

ABSTRACT

Cervical cancer was one of the global commitments in Sustainable Development Goals (SDGs). Cervical cancer represents 7.5% of all deaths caused by cancer in women. The method to screening cervical cancer are pap smear test and Visual Inspection of Acetic Acid (VIA) test. Indonesia is the second country in the world has the most cervical cancer cases. This study aims to determine the behavior of cervical cancer prevention on housewives. This is kuantitatif research with cross sectional design. The subjects of this study were housewives in the Kota Yogyakarta and Sleman regency amounting to 350 people. The analysis used is univariate, bivariate and multivariate analysis. The results showed that there were 45.1% of mothers' have did the implementation of VIA. There is a relationship of education ($p = 0.024$), level of knowledge ($p = 0.036$), and maternal attitudes with cervical cancer prevention behavior. There were no correlation between age ($p = 0$), parity ($p = 0.816$), and family income ($p = 0.174$) with the mother's behavior in the implementation of the VIA's test. Multivariate analysis showed maternal attitude ($CI = 0.335-0.788$) that affect the behavior of the VIA.

Copyright © 2018 Institute of Advanced Engineering and Science.
All rights reserved.

Corresponding Author:

Nanik Setiyawati,
Midwifery Department,
Helath Polytechnic Ministry of Helath Yogyakarta,
Mangkuyudan MJ III/304 Yogyakarta, Indonesia.
Email: nanikyogya@gmail.com

1. INTRODUCTION

Cervical cancer cases in 2012 included 528 000 new cases, which have been diagnosed worldwide and 85% occur in less developed areas. Cervical cancer causes 266 000 women to die of cervical cancer every year. Cervical cancer represents 7.5% of all deaths caused by cancer in women. From those data, we know that cervical cancer is a threat for the health that should be pressed down at least until 2030 which consist of curative action to the people living with cervical cancer [1].

Indonesia is the second country in the world after China has the most cervical cancer [1]. Based on data from the Ministry of Health in 2015, on average every hour the number of Cervical Cancer sufferers increased by 2.5 people and 1.1 women died of cervical cancer. The prevalence and estimation of cervical cancer patients in 2013 in Indonesia was 0.8 ‰ with an estimated total of 98,692 cases. Riau Islands Province, North Maluku Province, and D.I.Y Province (Yogyakarta Special Region) have the highest prevalence of cervical cancer which is 1.5 ‰ with a total of 2,703 cases [2].

Prevention of cervical cancer can be done by performing an early cervical health examination (screening), because symptoms of cervical cancer are not seen until the stage is more severe [3]. High-quality screening with cytology (Papanicolaou [Pap] testing) has markedly reduced mortality from squamous cell cervical cancer, which comprises 80% to 90% of cervical cancers [4-6]. Examination of early detection of cervical cancer with VIA is a visual examination of the cervix using vinegar, meaning seeing the cervix with the naked eye to detect abnormalities after applying acetic acid or vinegar (3-5%).

To improve the implementation of prevention and early detection of cancer in women in Indonesia, the government is optimizing the cervical cancer early detection program for the 2015-2019 period. The effort taken was an early detection movement through the method of simultaneous VIA examination in all parts of Indonesia on April 21, 2015. This movement will last for 5 years and it is expected that by 2019 the number of WUS (women of childbearing age) who are carried out early detection reaches 50 percent [2].

On the word of Hae Won Kim and Duck Hee Kim, 2015 that The mothers' awareness and preparedness with respect to the prevention of cervical cancer in their adolescent daughters were low and inadequate. Mothers should be informed and motivated to play a role in the education of their daughters regarding cervical cancer prevention. Strategies for disseminating information regarding early cervical cancer prevention for adolescent girls are recommended by communicating with both the girls and their mothers and providing them with education regarding cervical cancer prevention [8]. This study aims to determine the factors that influence mother in behavior of cervical cancer prevention.

2. RESEARCH METHOD

This study used analytic survey with cross-sectional design. This study conducted on housewives in the Kota Yogyakarta and Sleman who were members of the Family Empowerment and Welfare amounting to 350 people. This study was conducted in 2016. The population target in this study was housewives who were members of the Family Empowerment and Welfare to manage the confounding variables. To determine the sample size, we used the formula for sample size with calculation of proportion as per Lemeshow with minimum sample size obtained was 350 respondent. Proportional sampling was used to determine the sample chosen. Variables in this study used the theoretical framework approach of Theory of Precede-Proceed (Lawrence Green). Independent variable consist of the characteristic of respondent (age, parity, education, family income), knowledge level about VIA, and mother' attitude toward VIA. The dependent variable was cervical cancer prevention behavior in mother. This study took primary data. Instrument used in this study was questionnaire. Questionnaire used was development from questionnaire about mothers' knowledge, attitude and behavior toward VIA which have been tested the validity and reliability at Bantul district.

3. RESULTS AND ANALYSIS

This study conducted on 350 respondent was housewives were members of the Family Empowerment and Welfare. The characteristic of the respondent showed in Table 1. Table 1 shows that the major proportion of respondents were aged >35 years old (76.6%), with the highest parity was secundipara in the amount of 166 respondents (47.5%). More than half of the respondents with secondary education amounted to 191 respondents (54.6%). Family income >Rp. 1.450.000,00 as many as 189 respondents (54%).

Table 1. Respondent's Characteristic

Characteristic	Amount (n=350)	Percentage (%)
Age		
≤35 years	82	23.4
>35 years	268	76.6
Parity		
Nulipara	6	1.7
Primipara	83	23.7
Secundipara	166	47.5
Multi gravida	95	27.1
Education		
Basic	91	26.0
Medium	191	54.6
High	68	19.4
Family Income		
> Rp.1.450.000,00	189	54.0
≤ Rp.1.450.000,00	161	46.0

Table 2 shows that most respondent have not enough knowledge about VIA (80.6%). There are still 35 respondents (10%) with a lack of knowledge, the role of health workers is needed in providing insight to mothers regarding early detection of cervical cancer (VIA). Most respondent have positive attitude toward VIA (54.6%). The behavior of the VIA inspection there are 192 respondents (54.9%) for not VIA.

Table 2. Mothers' Knowledge Level, Attitude, and Behavior toward VIA Test

Characteristic	Amount (n=350)	Percentage (%)
Knowledge level about VIA		
Good	33	9.4
Adequate	282	80.6
Poor	35	10.0
Mothers' attitude toward VIA		
Positive	191	54.6
Negative	159	45.4
Mothers' behavior toward VIA		
Do VIA	158	45.1
Not Do VIA	192	54.9

Bivariate analysis was used to analyze independent and dependent variables. The relation between independent and dependent variables are showed in Table 3. Table 3 shows that factors relating to the behavior of VIA with a p-value <0.05 was educational p=0.024, p=0.036 level of knowledge, and attitudes toward VIA inspection p=0.002. 268 respondents with age >35 years as many as 147 respondents (54.9%) did not do VIA. 166 respondents with secundipara parity as many as 90 respondents (54.2%) did not do VIA. 189 respondents with family income >Rp 1.450.000,00 as many as 110 respondents (58.2%) did not do an VIA. So it can be concluded that there is no significant relationship between parity and family income with IVA behavior (p-value 0.816 and 0.174).

Table 3. The Relationship between Mothers' Characteristic, Knowledge Level, Attitude, and Behavior toward VIA

Variable	Behavior toward VIA				Total		p-value
	Not Do VIA n=192		Do VIA n=158		n=350		
	n	%	n	%	n	%	
Age							
≤35 years	45	54.9	37	45.1	82	100	
> 35 years	147	54.9	121	45.1	268	100	
Parity							
Nulipara	3	50	3	50	6	100	0.816
Primipara	48	57.8	35	42.2	83	100	
Sekundipara	90	54.2	76	45.8	166	100	
Multipara	51	53.7	44	46.3	95	100	
Education Level							
Basic	57	62.6	34	37.4	91	100	0.024*
Medium	102	53.4	89	46.6	191	100	
High	33	48.5	35	51.5	68	100	
Family Income							
> Rp.1.450.000,00	110	58.2	79	41.8	189	100	0.174
≤Rp.1.450.000,00	82	51	79	49	161	100	
Knowledge level							
Superior	16	48.5	17	51.5	33	100	0.036*
Satisfactory	150	53.2	132	46.8	282	100	
Unsatisfactory	26	74.3	9	25.7	35	100	
Attitude toward VIA							
Positive	119	62.3	72	37.7	191	100	0.002*
Negative	73	45.9	86	54.1	159	100	

Multivariate analysis were done to independent variables that have p <0.002 altogether. The result of multivariate analysis showed in Table 4. Table 4 shows that the results of statistical tests with logistic regression found that the attitude has a p-value of 0.002. This shows that the factor that most influences the behavior toward VIA is the attitude of respondents. Its has a p-value of 0.002>0.05.

Table 4. Multivariate Analysis Result

	B	Sig.	Exp(B)	95.0% C.I. for EXP(B)	
				Lower	Upper
Attitude toward VIA test	-.666	.002	.514	.335	788
Constant	.164	.303	1.178		

Cervical cancer could be prevented by screening VIA. The results of abudukadeer A et al showed that women between the age group 31 to 40 years had more awareness about cervical cancer [9]. Respondents older than 30 years have a possibility of having a precancerous lesion is higher than under 30 years of age so that respondents in that age will find it more important to conduct tests VIA [10]. The results of this study found no correlation of age to the prevention behavior of IVA cervical cancer ($p=0$). The results of this study are in accordance with Gustiana D's study that there was no significant relationship between age and early detection of cervical cancer $p=0.306$. It can be associated with susceptibility to disease. Age can not be used as a benchmark for a person to prevent cervical cancer. This can be due to ignorance, no complaints, nor do to prevent cervical cancer has not been necessary [11].

The results of previous studies conducted by Yuliwati also showed that there was no significant relationship between age and maternal behavior to check VIA [12]. However, different research results Eva who showed that age has a significant relationship with the knowledge that motivates behavior of women for the early detection of cervical cancer by VIA in the district of Central Bogor $p=0.001$ [13].

The results of Heni P et al's research showed that parity had a significant relationship to precancerous cervical lesions [14]. While the chi square test results in this study indicate that there is no significant relationship between parity against the IVA behavior $p=0.816$. The results of this study are inconsistent with Sri Dinengsih research and Erry Sitanggang states that there is a significant relationship between parity with IVA inspection. A third parity or more at risk of developing cervical cancer. Multiparity suspected of causing a decrease in endurance. So it is necessary to do the IVA method to detect early cervical cancer [15].

The level of education a person can support or influence a person's level of knowledge and a low level of education is always related to information and knowledge is limited, the higher one's education the higher one's own understanding and knowledge of information obtained would be even higher [16]. Based on the results of this study indicate that there is a significant relationship between education with VIA behavior $p=0.024$. This research is in line with research conducted by Yao Jia on several women in the city of Wufeng, China who explained that education affects the implementation of early detection of cervical cancer by 0.00 ($p=0.05$) with the most characteristics at the low level of education (48.4%) who are willing to carry out early detection of cervical cancer [17]. This is also confirmed by the results of further research by Eva, explains that the education factor has a significant relationship with the level of knowledge possessed by women of childbearing age, where the group of higher education have a proportion of knowledge both larger than the group of secondary education and low [13]. The level of education is one of the factors that determine a person's knowledge and perception of the importance of a case, including the importance of early detection of cervical cancer, caused by highly educated person will be broader view and more receptive to the ideas and new ways of life. It can be concluded that highly educated people will make early detection of cervical cancer [18].

The results showed that no significant relationship between family income on behavior IVA $p=0.174$. The results are consistent with research Gustiana states that there is no relationship to the economic status of cervical cancer prevention behavior ($p=0.561$), research found that respondents with lower economic status also have a good preventive health behaviors [11]. Another study conducted by Wahyuni states that there is no economic status relationship to the behavior of early detection of cervical cancer [19]. Socio-economic conditions affecting the health status change process as it affects the thoughts or beliefs that can lead to changes in health behavior. The incidence of cervical cancer is twice as large in women who have low social classes. In contrast to the results of research Febriani CA Women who have high incomes likely better prevention of cervical cancer, compared to women with low incomes [17].

Increased knowledge can change people's behavior from negative to positive, in addition to the knowledge also build trust [11]. Having knowledge of a disease, the concept of a disease will be formed in the individual so that it will determine a person's health behavior [20]. Lack of related knowledge about cervical cancer can be an important factor for the high incidence of cervical cancer [9]. Based on the results of chi-square test showed that there is a relationship with the level of knowledge of the behavior of IVA $p=0.036$. The results are consistent with research conducted by Yao Jia on women in the town of Wufeng China stated there is a significant relationship between knowledge with behavioral examination early detection of cervical cancer through IVA method with $p=0.000$ [17]. In addition, research conducted by Nesrin, Turkish women gained knowledge influence the early detection of cervical cancer ($p=0.001$) [21]. Based on research conducted by Khosidah, good knowledge possessed by women of childbearing age about cervical cancer and its examination can be a motivating factor to try to avoid cervical cancer. Various sources of information that can now be easily accessed by women of childbearing age allow knowledge of cervical cancer and the examination will be better. Lack of knowledge and awareness of the importance of examination is an inhibiting factor for screening for cervical cancer [22].

Attitude is one of the personality elements that a person has to determine his actions and behave towards an object accompanied by positive and negative feelings. Attitudes influence the formation of interest because of the tendency in the subject to accept or reject an object that is good or not [23]. The chi square test

results show that there is a significant relationship between attitudes toward behavior IVA $p=0.002$. The results of this study are in line with Febriani CA's research the attitude of adult women in having a significant relationship with the examination of early detection of cervical cancer $p=0.025$ [18]. TRA explains that beliefs can affect attitudes and social norms which would change the form of the desire well behaved guided or happen in an individual's behavior. This theory confirms the role of one's intention in determining whether a behavior will occur [24].

4. CONCLUSION

Respondent joined this study were mostly aged >35 years old, secundipara, medium level of knowledge, family income more than minimum income regional. There is a relationship of education, level of knowledge, and attitudes with cervical cancer prevention behavior. There is no correlation between age, parity, and family income with the mother's behavior on VIA tests. Attitude towards VIA tests is the most affect to behavior of the VIA.

ACKNOWLEDGEMENTS

We would like to thank the women and research assistants who participated in this study. The study was funded by Health Polytechnic Ministry of Health Yogyakarta, Indonesia.


REFERENCES

- [1] World Health Organization (WHO). Care of the preterm and low-birth-weight newborn world prematurity day - " Let Them Thrive ." 2017;1-5.
- [2] Septarini NW, Kurniati DPY, Wiryanthini IAD, Putra AWGAE, Sutarga LM. Inisiasi kader desa peduli kesehatan reproduksi (kdpkr) serta deteksi dini kanker leher rahim di desa pengotan kabupaten bangli bali. *Bul Udayana Mengabdikan*. 2017;16(September):107-15.
- [3] R I, Kementerian Kesehatan. Situasi penyakit kanker. *InfoDATIN* (2015):6-9.
- [4] Nordianti ME, Wahyono B. Determinan kunjungan inspeksi visual asam asetat di puskesmas kota semarang. *HIGEIA J PUBLIC Heal Res Dev*. 2018;2(1):33-44.
- [5] Gustafsson L, Ponten J, Bergstrom R, Adami HO. International incidence rates of invasive cervical cancer before cytological screening. *Int J cancer*. 1997;71: 159-165.
- [6] Gustafsson L, Ponten J, Zack M, Adami HO. International incidence rates of invasive cervical cancer after introduction of cytological screening. *Cancer Causes Control*. 1997;8:755-763.
- [7] Parkin DM, Bray F. Chapter 2: The burden of HPV-related cancers. *accine*. 2006; 24(suppl 3):S3/11-25.
- [8] Gustafsson L, Ponten J, Bergstrom R, Adami HO. International incidence rates of invasive cervical cancer before cytological screening. *Int J Cancer*. 1997;71:159-165.
- [9] Kim HW, Kim DH. Awareness of cervical cancer prevention among mothers of adolescent daughters in korea : qualitative research. *BMJ Open*. 2015;1-10.
- [10] Abudukadeer A, Azam S, Mutailipu AZ, Qun L, Guilin G, Mijiti S. Knowledge and attitude of Uyghur women in Xinjiang province of China related to the prevention and early detection of cervical cancer. *World J Surg Oncol*. 2015;13(110):1-7.
- [11] Mulyati S, Suwarsa O, D A IF. Pengaruh media film terhadap sikap ibu pada deteksi dini kanker serviks. *J Kesehat Masy*. 2015;11(2):16-24.
- [12] Gustiana D, Dewi YI, Nurchayati S, Studi P, Keperawatan I, Riau U. Faktor-faktor yang berhubungan dengan perilaku pencegahan kanker serviks pada wanita usia subur. *JOM PSIK*. 2014;1(2):1-8.
- [13] Yuliwati. Faktor-faktor yang berhubungan dengan perilaku wus dalam deteksi dini kanker leher rahim metode IVA Di Wilayah Puskesmas Prembun 2012. 2012; Available from: <http://repository.ui.ac.id>.
- [14] Sulistiawati E. Pengetahuan tentang faktor risiko, perilaku dan deteksi dini kanker serviks dengan inspeksi visual asam asetat (IVA) pada wanita di kecamatan bogor tengah. *Bogor. Bul Penelit Kesehat*. 2014;42(3):192-202.
- [15] Purwaningsih H, Pradjatno H. Risk factors associated with the occurrence of precancerous cervical lesions at health centres of Karanganyar regency. *Int J Community Med and Public Heal*. 2016;3(6):1437-41.
- [16] Dinengsih S, Sitanggang E. Analisis faktor perilaku deteksi dini kanker serviks dengan metode IVA (Inspeksi Visual Asam Asetat). *URECOL*. 2018;37-45.
- [17] Parapat FT, S HS, Sc M, Saraswati LD, Epid M. Faktor faktor yang berhubungan dengan perilaku deteksi dini kanker leher rahim metode inspeksi visual asam asetat di puskesmas candiroto kabupaten temanggung. *J Kesehat Masy*. 2016;4(4):363-70.
- [18] Yao J. Knowledge about cervical cancer and barriers of screening program among women in wufeng county, a high-incidence region of cervical cancer in china. 2013.
- [19] Febriani CA. Faktor-faktor yang berhubungan dengan deteksi dini kanker leher rahim di kecamatan gisting kabupaten tanggamus lampung. *J Kesehat*. 2015;VII(2):228-37.
- [20] Wahyuni S. Faktor-faktor yang mempengaruhi perilaku deteksi dini kanker serviks di kecamatan ngampel kabupaten kendal jawa tengah. 2013. Available from: <https://bem.unimus.ac.id>.

Commented [WU2]: No 7 dan 25 tidak dirujuk/sitasi di body text

- [21] Situmorang. Hubungan pengetahuan dan sikap dengan perilaku deteksi dini pada penderita kanker serviks di RSUD dr. Kariadi Semarang Tahun 2015. *J Kesehatan Masy.* 2016;4(1):76–82.
- [22] Reis N. Knowledge, behavior, and beliefs for cervical cancer and screening in turkey. 2012; Available from: doi:<http://dx.doi.org/10.7314/APJCP.2012.13.4.1463>
- [23] Khosidah A. Faktor-faktor yang mempengaruhi ibu rumah tangga dalam melakukan tes IVA sebagai upaya deteksi dini kanker serviks. *J Ilm Kebidanan.* 2015;6(2):94–105.
- [24] Sholihah A. Hubungan antara sikap pencegahan kanker serviks dengan minat deteksi dini menggunakan IVA pada WUS di desa mojolaban sukoharjo. *Artik Ilm.* 2015;20(1).
- [25] Green LW. *Health promotion planning an educational and environmental approach.* Mountain View Toronto - London: Mayfield Publishing Company; 2000.

7. BUKTI KONFIRMASI HASIL REVIEW KETIGA DAN BUKTI REVISI HASIL REVIEW KETIGA (10 NOVEMBER 2018)



Prevention of Cervical Cancer on Mother in Yogyakarta, Indonesia

Nanik Setiyawati, Niken Meilani
Poltekkes Kemenkes Yogyakarta

Article Info

Article history:

Received Sep 28, 2018

Revised Nov 8, 2018

Accepted Nov 15, 2018

Keyword:

Attitudes
Housewife
VIA test

ABSTRACT

Cervical cancer was one of the global commitments in Sustainable Development Goals (SDGs). Cervical cancer represents 7.5% of all deaths caused by cancer in women. The method to screening cervical cancer are pap smear test and Visual Inspection of Acetic Acid (VIA) test. Indonesia is the second country in the world has the most cervical cancer cases. This study aims to determine the behavior of cervical cancer prevention on housewives. This is kuantitatif research with cross sectional design. The subjects of this study were housewives in the Kota Yogyakarta and Sleman regency amounting to 350 people. The analysis used is univariate, bivariate and multivariate analysis. The results showed that there were 45.1% of mothers' have did the implementation of VIA. There is a relationship of education ($p = 0.024$), level of knowledge ($p = 0.036$), and maternal attitudes with cervical cancer prevention behavior. There were no correlation between age ($p = 0$), parity ($p = 0.816$), and family income ($p = 0.174$) with the mother's behavior in the implementation of the VIA's test. Multivariate analysis showed maternal attitude ($CI = 0.335-0.788$) that affect the behavior of the VIA.

Copyright © 2018 Institute of Advanced Engineering and Science.
All rights reserved.

Corresponding Author:

Nanik Setiyawati,
Midwifery Department,
Poltekkes Kemenkes Yogyakarta,
Mangkuyudan MJ III/304 Yogyakarta, Indonesia.
Email: nanikyogya@gmail.com

1. INTRODUCTION

Cervical cancer cases in 2012 included 528 000 new cases, which have been diagnosed worldwide and 85% occur in less developed areas. Cervical cancer causes 266 000 women to die of cervical cancer every year. Cervical cancer represents 7.5% of all deaths caused by cancer in women. From those data, we know that cervical cancer is a threat for the health that should be pressed down at least until 2030 which consist of curative action to the people living with cervical cancer [1].

Indonesia is the second country in the world after China has the most cervical cancer [1]. Based on data from the Ministry of Health in 2015, on average every hour the number of Cervical Cancer sufferers increased by 2.5 people and 1.1 women died of cervical cancer. The prevalence and estimation of cervical cancer patients in 2013 in Indonesia was 0.8 ‰ with an estimated total of 98,692 cases. Riau Islands Province, North Maluku Province, and D.I.Y Province (Yogyakarta Special Region) have the highest prevalence of cervical cancer which is 1.5 ‰ with a total of 2,703 cases [2].

Prevention of cervical cancer can be done by performing an early cervical health examination (screening), because symptoms of cervical cancer are not seen until the stage is more severe [3]. High-quality screening with cytology (Papanicolaou [Pap] testing) has markedly reduced mortality from squamous cell cervical cancer, which comprises 80% to 90% of cervical cancers [4-6]. Examination of early detection of cervical cancer with VIA is a visual examination of the cervix using vinegar, meaning seeing the cervix with the naked eye to detect abnormalities after applying acetic acid or vinegar (3-5%).

To improve the implementation of prevention and early detection of cancer in women in Indonesia, the government is optimizing the cervical cancer early detection program for the 2015-2019 period. The effort taken was an early detection movement through the method of simultaneous VIA examination in all parts of Indonesia on April 21, 2015. This movement will last for 5 years and it is expected that by 2019 the number of WUS (women of childbearing age) who are carried out early detection reaches 50 percent [2].

On the word of Hae Won Kim and Duck Hee Kim, 2015 that The mothers' awareness and preparedness with respect to the prevention of cervical cancer in their adolescent daughters were low and inadequate. Mothers should be informed and motivated to play a role in the education of their daughters regarding cervical cancer prevention. Strategies for disseminating information regarding early cervical cancer prevention for adolescent girls are recommended by communicating with both the girls and their mothers and providing them with education regarding cervical cancer prevention [7]. This study aims to determine the factors that influence mother in behavior of cervical cancer prevention.

2. RESEARCH METHOD

This study used analytic survey with cross-sectional design. This study conducted on housewives in the Kota Yogyakarta and Sleman who were members of the Family Empowerment and Welfare amounting to 350 people. This study was conducted in 2016. The population target in this study was housewives who were members of the Family Empowerment and Welfare to manage the confounding variables. To determine the sample size, we used the formula for sample size with calculation of proportion as per Lemeshow with minimum sample size obtained was 350 respondent. Proportional sampling was used to determine the sample chosen. Variables in this study used the theoretical framework approach of Theory of Precede-Proceed (Lawrence Green) [8]. Independent variable consist of the characteristic of respondent (age, parity, education, family income), knowledge level about VIA, and mother' attitude toward VIA. The dependent variable was cervical cancer prevention behavior in mother. This study took primary data. Instrument used in this study was questionnaire. Questionnaire used was development from questionnaire about mothers' knowledge, attitude and behavior toward VIA which have been tested the validity and reliability at Bantul district.

3. RESULTS AND ANALYSIS

This study conducted on 350 respondent was housewives were members of the Family Empowerment and Welfare. The characteristic of the respondent showed in Table 1. Table 1 shows that the major proportion of respondents were aged >35 years old (76.6%), with the highest parity was secundipara in the amount of 166 respondents (47.5%). More than half of the respondents with secondary education amounted to 191 respondents (54.6%). Family income >Rp. 1.450.000,00 as many as 189 respondents (54%).

Table 1. Respondent's Characteristic

Characteristic	Amount (n=350)	Percentage (%)
Age		
≤35 years	82	23.4
>35 years	268	76.6
Parity		
Nulipara	6	1.7
Primipara	83	23.7
Secundipara	166	47.5
Multi gravida	95	27.1
Education		
Basic	91	26.0
Medium	191	54.6
High	68	19.4
Family Income		
> Rp.1.450.000,00	189	54.0
≤Rp.1.450.000,00	161	46.0

Table 2 shows that most respondent have not enough knowledge about VIA (80.6%). There are still 35 respondents (10%) with a lack of knowledge, the role of health workers is needed in providing insight to mothers regarding early detection of cervical cancer (VIA). Most respondent have positive attitude toward VIA (54.6%). The behavior of the VIA inspection there are 192 respondents (54.9%) for not VIA.

Table 2. Mothers' Knowledge Level, Attitude, and Behavior toward VIA Test

Characteristic	Amount (n=350)	Percentage (%)
Knowledge level about VIA		
Good	33	9.4
Adequate	282	80.6
Poor	35	10.0
Mothers' attitude toward VIA		
Positive	191	54.6
Negative	159	45.4
Mothers' behavior toward VIA		
Do VIA	158	45.1
Not Do VIA	192	54.9

Bivariate analysis was used to analyze independent and dependent variables. The relation between independent and dependent variables are showed in Table 3. Table 3 shows that factors relating to the behavior of VIA with a p-value <0.05 was educational p=0.024, p=0.036 level of knowledge, and attitudes toward VIA inspection p=0.002. 268 respondents with age>35 years as many as 147 respondents (54.9%) did not do VIA. 166 respondents with secundipara parity as many as 90 respondents (54.2%) did not do VIA. 189 respondents with family income >Rp 1.450.000,00 as many as 110 respondents (58.2%) did not do an VIA. So it can be concluded that there is no significant relationship between parity and family income with IVA behavior (p-value 0.816 and 0.174).

Table 3. The Relationship between Mothers' Characteristic, Knowledge Level, Attitude, and Behavior toward VIA

Variable	Behavior toward VIA				p-value	
	Not Do VIA n=192		Do VIA n=158		Total n=350	%
Age						
≤35 years	45	54.9	37	45.1	82	100
> 35 years	147	54.9	121	45.1	268	100
Parity						
Nulipara	3	50	3	50	6	100
Primipara	48	57.8	35	42.2	83	100
Sekundipara	90	54.2	76	45.8	166	100
Multipara	51	53.7	44	46.3	95	100
Education Level						
Basic	57	62.6	34	37.4	91	100
Medium	102	53.4	89	46.6	191	100
High	33	48.5	35	51.5	68	100
Family Income						
> Rp.1.450.000,00	110	58.2	79	41.8	189	100
≤Rp.1.450.000,00	82	51	79	49	161	100
Knowledge level						
Superior	16	48.5	17	51.5	33	100
Satisfactory	150	53.2	132	46.8	282	100
Unsatisfactory	26	74.3	9	25.7	35	100
Attitude toward VIA						
Positive	119	62.3	72	37.7	191	100
Negative	73	45.9	86	54.1	159	100

Multivariate analysis were done to independent variables that have p <0.002 altogether. The result of multivariate analysis showed in Table 4. Table 4 shows that the results of statistical tests with logistic regression found that the attitude has a p-value of 0.002. This shows that the factor that most influences the behavior toward VIA is the attitude of respondents. Its has a p-value of 0.002>0.05.

Table 4. Multivariate Analysis Result

	B	Sig.	Exp(B)	95.0% C.I. for EXP(B)	
				Lower	Upper
Attitude toward VIA test	-.666	.002	.514	.335	788
Constant	.164	.303	1.178		

Cervical cancer could be prevented by screening VIA. The results of abudukadeer A et al showed that women between the age group 31 to 40 years had more awareness about cervical cancer [9]. Respondents older than 30 years have a possibility of having a precancerous lesion is higher than under 30 years of age so that respondents in that age will find it more important to conduct tests VIA [10]. The results of this study found no correlation of age to the prevention behavior of IVA cervical cancer ($p=0$). The results of this study are in accordance with Gustiana D's study that there was no significant relationship between age and early detection of cervical cancer $p=0.306$. It can be associated with susceptibility to disease. Age can not be used as a benchmark for a person to prevent cervical cancer. This can be due to ignorance, no complaints, nor do to prevent cervical cancer has not been necessary [11].

The results of previous studies conducted by Yuliwati also showed that there was no significant relationship between age and maternal behavior to check VIA [12]. However, different research results Eva who showed that age has a significant relationship with the knowledge that motivates behavior of women for the early detection of cervical cancer by VIA in the district of Central Bogor $p=0.001$ [13].

The results of Heni P et al's research showed that parity had a significant relationship to precancerous cervical lesions [14]. While the chi square test results in this study indicate that there is no significant relationship between parity against the IVA behavior $p=0.816$. The results of this study are inconsistent with Sri Dinengsih research and Erry Sitanggang states that there is a significant relationship between parity with IVA inspection. A third parity or more at risk of developing cervical cancer. Multiparity suspected of causing a decrease in endurance. So it is necessary to do the IVA method to detect early cervical cancer [15].

The level of education a person can support or influence a person's level of knowledge and a low level of education is always related to information and knowledge is limited, the higher one's education the higher one's own understanding and knowledge of information obtained would be even higher [16]. Based on the results of this study indicate that there is a significant relationship between education with VIA behavior $p=0.024$. This research is in line with research conducted by Yao Jia on several women in the city of Wufeng, China who explained that education affects the implementation of early detection of cervical cancer by 0.00 ($p=0.05$) with the most characteristics at the low level of education (48.4%) who are willing to carry out early detection of cervical cancer [17]. This is also confirmed by the results of further research by Eva, explains that the education factor has a significant relationship with the level of knowledge possessed by women of childbearing age, where the group of higher education have a proportion of knowledge both larger than the group of secondary education and low [13]. The level of education is one of the factors that determine a person's knowledge and perception of the importance of a case, including the importance of early detection of cervical cancer, caused by highly educated person will be broader view and more receptive to the ideas and new ways of life. It can be concluded that highly educated people will make early detection of cervical cancer [18].

The results showed that no significant relationship between family income on behavior IVA $p=0.174$. The results are consistent with research Gustiana states that there is no relationship to the economic status of cervical cancer prevention behavior ($p=0.561$), research found that respondents with lower economic status also have a good preventive health behaviors [11]. Another study conducted by Wahyuni states that there is no economic status relationship to the behavior of early detection of cervical cancer [19]. Socio-economic conditions affecting the health status change process as it affects the thoughts or beliefs that can lead to changes in health behavior. The incidence of cervical cancer is twice as large in women who have low social classes. In contrast to the results of research Febriani CA Women who have high incomes likely better prevention of cervical cancer, compared to women with low incomes [17].

Increased knowledge can change people's behavior from negative to positive, in addition to the knowledge also build trust [11]. Having knowledge of a disease, the concept of a disease will be formed in the individual so that it will determine a person's health behavior [20]. Lack of related knowledge about cervical cancer can be an important factor for the high incidence of cervical cancer [9]. Based on the results of chi-square test showed that there is a relationship with the level of knowledge of the behavior of IVA $p=0.036$. The results are consistent with research conducted by Yao Jia on women in the town of Wufeng China stated there is a significant relationship between knowledge with behavioral examination early detection of cervical cancer through IVA method with $p=0.000$ [17]. In addition, research conducted by Nesrin, Turkish women gained knowledge influence the early detection of cervical cancer ($p=0.001$) [21]. Based on research conducted by Khosidah, good knowledge possessed by women of childbearing age about cervical cancer and its examination can be a motivating factor to try to avoid cervical cancer. Various sources of information that can now be easily accessed by women of childbearing age allow knowledge of cervical cancer and the examination will be better. Lack of knowledge and awareness of the importance of examination is an inhibiting factor for screening for cervical cancer [22].

Attitude is one of the personality elements that a person has to determine his actions and behave towards an object accompanied by positive and negative feelings. Attitudes influence the formation of interest because of the tendency in the subject to accept or reject an object that is good or not [23]. The chi square test results show that there is a significant relationship between attitudes toward behavior IVA $p=0.002$. The results of this study are in line with Febriani CA's research the attitude of adult women in having a significant relationship with the examination of early detection of cervical cancer $p=0.025$ [18]. TRA explains that beliefs can affect attitudes and social norms which would change the form of the desire well behaved guided or happen in an individual's behavior. This theory confirms the role of one's intention in determining whether a behavior will occur [24].

4. CONCLUSION

Respondent joined this study were mostly aged >35 years old, secundipara, medium level of knowledge, family income more than minimum income regional. There is a relationship of education, level of knowledge, and attitudes with cervical cancer prevention behavior. There is no correlation between age, parity, and family income with the mother's behavior on VIA tests. Attitudes towards VIA tests is the most affect to behavior of the VIA.

ACKNOWLEDGEMENTS

We would like to thank the women and research assistants who participated in this study. The study was funded by Health Polytechnic Ministry of Health Yogyakarta, Indonesia.

REFERENCES

- [1] World Health Organization (WHO). Care of the preterm and low-birth-weight newborn world prematurity day - "Let Them Thrive ." 2017;1-5.
- [2] Septarini NW, Kurniati DPY, Wiryanthini IAD, Putra AWGAE, Sutarga LM. Inisiasi kader desa peduli kesehatan reproduksi (kdpkr) serta deteksi dini kanker leher rahim didesa pengotan kabupaten bangli bali. *Bul Udayana Mengabdi*. 2017;16(September):107-15.
- [3] R I, Kementerian Kesehatan. Situasi penyakit kanker. *InfoDATIN* (2015):6-9.
- [4] Nordianti ME, Wahyono B. Determinan kunjungan inspeksi visual asam asetat di puskesmas kota semarang. *HIGEIA J PUBLIC Heal Res Dev*. 2018;2(1):33-44.
- [5] Gustafsson L, Ponten J, Bergstrom R, Adami HO. International incidence rates of invasive cervical cancer before cytological screening. *Int J Cancer*. 1997;71:159-165.
- [6] Gustafsson L, Ponten J, Zack M, Adami HO. International incidence rates of invasive cervical cancer after introduction of cytological screening. *Cancer Causes Control*. 1997;8:755-763.
- [7] Gustafsson L, Ponten J, Bergstrom R, Adami HO. International incidence rates of invasive cervical cancer before cytological screening. *Int J Cancer*. 1997;71:159-165.
- [8] Green LW. Health promotion planning an educational and environmental approach. Mountain View Toronto - London: Mayfield Publishing Company; 2000.
- [9] Kim HW, Kim DH. Awareness of cervical cancer prevention among mothers of adolescent daughters in Korea: qualitative research. *BMJ Open*. 2015;1-10.
- [10] Abudukadeer A, Azam S, Mutailipu AZ, Qun L, Guilin G, Mijiti S. Knowledge and attitude of Uyghur women in Xinjiang province of China related to the prevention and early detection of cervical cancer. *World J Surg Oncol*. 2015;13(110):1-7.
- [11] Mulyati S, Suwarsa O, D A IF. Pengaruh media film terhadap sikap ibu pada deteksi dini kanker serviks. *J Kesehat Masy*. 2015;11(2):16-24.
- [12] Gustiana D, Dewi YI, Nurchayati S, Studi P, Keperawatan I, Riau U. Faktor-faktor yang berhubungan dengan perilaku pencegahan kanker serviks pada wanita usia subur. *JOM PSIK*. 2014;1(2):1-8.
- [13] Yuliwati. Faktor-faktor yang berhubungan dengan perilaku wus dalam deteksi dini kanker leher rahim metode IVA Di Wilayah Puskesmas Prembun 2012. 2012; Available from: <http://repository.ui.ac.id>.
- [14] Sulistiowati E. Pengetahuan tentang faktor risiko, perilaku dan deteksi dini kanker serviks dengan inspeksi visual asam asetat (IVA) pada wanita di kecamatan bogor tengah. Bogor. *Bul Penelit Kesehat*. 2014;42(3):192-202.
- [15] Purwaningsih H, Prajatmo H. Risk factors associated with the occurrence of precancerous cervical lesions at health centres of Karanganyar regency. *Int J Community Med and Public Heal*. 2016;3(6):1437-41.
- [16] Dinengsih S, Sitanggang E. Analisis faktor perilaku deteksi dini kanker serviks dengan metode IVA (Inspeksi Visual Asam Acetat). *URECOL*. 2018;37-45.
- [17] Parapat FT, S HS, Sc M, Saraswati LD, Epid M. Faktor faktor yang berhubungan dengan perilaku deteksi dini kanker leher rahim metode inspeksi visual asam asetat di puskesmas candirot kabupaten temanggung. *J Kesehat Masy*. 2016;4(4):363-70.
- [18] Yao J. Knowledge about cervical cancer and barriers of screening program among women in wufeng county, a high-incidence region of cervical cancer in china. 2013.

Commented [WU2]: No 7 dan 25 tidakdirujuk/sitasi di body text
Jawab. Terima kasih.
Untuk No 7 kami delete dan ternyata nomor 25 adalah nomor 8. Dan yang nomor 8 seharusnya 7. Terima kasih

- [19] Febriani CA. Faktor-faktor yang berhubungan dengan deteksi dini kanker leher rahim di kecamatan gisting kabupaten tanggamus lampung. *J Kesehat*. 2015;VII(2):228–37.
- [20] Wahyuni S. Faktor-faktor yang mempengaruhi perilaku deteksi dini kanker serviks di kecamatan ngampel kabupaten kendal jawa tengah. 2013. Available from: <https://bem.unimus.ac.id>.
- [21] Situmorang. Hubungan pengetahuan dan sikap dengan perilaku deteksi dini pada penderita kanker serviks di RSUP dr. Kariadi semarang Tahun 2015. *J Kesehat Masy*. 2016;4(1):76–82.
- [22] Reis N. Knowledge, behavior, and beliefs for cervical cancer and screening in turkey. 2012; Available from: [doi:http://dx.doi.org/10.7314/APJCP.2012.13.4.1463](http://dx.doi.org/10.7314/APJCP.2012.13.4.1463)
- [23] Khosidah A. Faktor-faktor yang mempengaruhi ibu rumah tangga dalam melakukan tes IVA sebagai upaya deteksi dini kanker serviks. *J Ilm Kebidanan*. 2015;6(2):94–105.
- [24] Sholihah A. Hubungan antara sikap pencegahan kanker serviks dengan minat deteksi dini menggunakan IVA pada WUS di desa mojolaban sukoharjo. *Artik Ilm*. 2015;20(1).

ETHICAL CLEARANCE

Keterangan:

Judul EC: Penelitian Determinan Sikap Rumah Tangga terhadap tes HIV dan AIDS di Yogyakarta.

Dalam penelitian tersebut juga dikaji tentang perilaku Pap Smear.



KOMISI ETIK PENELITIAN KESEHATAN POLITEKNIK KESEHATAN KEMENKES YOGYAKARTA

Jl. Tatabumi No.3 Banyuraden, Gamping, Sleman, D.I.Yogyakarta Telp/Fax. 0274-617601

Website : www.komisi-etik.poltekkesjogja.ac.id | Email : komisietik.poltekkesjogja@gmail.com



PERSETUJUAN KOMISI ETIK No. LB.01.01/KE/XXI/099/2016

Judul	:	Determinan Sikap Ibu Rumah Tangga terhadap Tes HIV dan AIDS di Yogyakarta
Dokumen	:	1. Protokol 2. Persetujuan Setelah Penjelasan (PSP) 3. <i>Informed consent</i>
Nama Peneliti	:	Nanik Setiyawati, SST, S.Pd, M.Kes
Dokter/ Ahli medis yang bertanggungjawab	:	-
Tanggal Kelaikan Etik	:	2 Mei 2016
Institusi peneliti	:	Poltekkes Kemenkes Yogyakarta


Komisi Etik Penelitian Kesehatan (KEPK) Politeknik Kesehatan Kementerian Kesehatan Yogyakarta menyatakan bahwa protokol diatas telah memenuhi prinsip etis berdasarkan pada Deklarasi Helsinki 1975 dan oleh karena itu penelitian tersebut dapat dilaksanakan.

Surat Kelaikan Etik ini berlaku 1 (satu) tahun sejak tanggal terbit.

Komisi Etik Penelitian Kesehatan (KEPK) Politeknik Kesehatan Kementerian Kesehatan Yogyakarta memiliki hak untuk memantau kegiatan penelitian setiap saat. Peneliti wajib menyampaikan laporan akhir setelah penelitian selesai atau laporan kemajuan penelitian jika dibutuhkan.

Demikian. surat ini dibuat untuk dapat dipergunakan sebagaimana mestinya.

Ketua



Joko Susilo, SKM., M.Kes
Ketua KEPK Poltekkes Kemenkes Yogyakarta

TURNITIN

10._IJPHS_Desember_2018_P2.p df *by*

Submission date: 03-Oct-2021 10:32AM (UTC-0500)

Submission ID: 1663841435

File name: 10._IJPHS_Desember_2018_P2.pdf (211.65K)

Word count: 3731

Character count: 19888

Prevention of Cervical Cancer among Mother in Yogyakarta, Indonesia

Nanik Setiyawati, Niken Meilani
Poltekkes Kemenkes Yogyakarta, Indonesia

Article Info

Article history:

Received Sep 28, 2018

Revised Nov 8, 2018

Accepted Nov 23, 2018

Keyword:

Attitudes

Cervical cancer

Housewife

VIA test

ABSTRACT

Cervical cancer was one of the global commitments in Sustainable Development Goals (SDGs). Cervical cancer represents 7.5% of all deaths caused by cancer in women. The method to screening cervical cancer is pap smear test and Visual Inspection of Acetic Acid (VIA) test. Indonesia is the second country in the world has the most cervical cancer cases. This study aims to determine the behavior of cervical cancer prevention on housewives. This is kuantitatif research with cross sectional design. The subjects of this study were housewives in the Kota Yogyakarta and Sleman regency amounting to 350 people. The analysis used is univariate, bivariate and multivariate analysis. The results showed that there were 45.1% of mothers' have did the implementation of VIA. There is a relationship of education ($p=0.024$), level of knowledge ($p=0.036$), and maternal attitudes with cervical cancer prevention behavior. There were no correlation between age ($p=0$), parity ($p=0.816$), and family income ($p=0.174$) with the mother's behavior in the implementation of the VIA's test. Multivariate analysis showed maternal attitude ($CI=0.335-0.788$) that affect the behavior of the VIA.

Copyright © 2018 Institute of Advanced Engineering and Science.
All rights reserved.

Corresponding Author:

Nanik Setiyawati,

Midwifery Department,

Poltekkes Kemenkes Yogyakarta,

Mangkuyudan MJ III/304 Yogyakarta, Indonesia.

Email: nanikyogya@gmail.com

1. INTRODUCTION

Cervical cancer cases in 2012 included 528 000 new cases, which have been diagnosed worldwide and 85% occur in less developed areas. Cervical cancer causes 266 000 women to die of cervical cancer every year. Cervical cancer represents 7.5% of all deaths caused by cancer in women. From those data, we know that cervical cancer is a threat for the health that should be pressed down at least until 2030 which consist of curative action to the people living with cervical cancer [1].

Indonesia is the second country in the world after China has the most cervical cancer [1]. Based on data from the Ministry of Health in 2015, on average every hour the number of Cervical Cancer sufferers increased by 2.5 people and 1.1 women died of cervical cancer. The prevalence and estimation of cervical cancer patients in 2013 in Indonesia was 0.8 % with an estimated total of 98,692 cases. Riau Islands Province, North Maluku Province, and D.I.Y Province (Yogyakarta Special Region) have the highest prevalence of cervical cancer which is 1.5% with a total of 2,703 cases [2].

Prevention of cervical cancer can be done by performing an early cervical health examination (screening), because symptoms of cervical cancer are not seen until the stage is more severe [3]. High-quality screening with cytology (Papanicolaou [Pap] testing) has markedly reduced mortality from squamous cell cervical cancer, which comprises 80% to 90% of cervical cancers [4-6]. Examination of early detection of

cervical cancer with VIA is a visual examination of the cervix using vinegar, meaning seeing the cervix with the naked eye to detect abnormalities after applying acetic acid or vinegar (3-5%).

To improve the implementation of prevention and early detection of cancer in women in Indonesia, the government is optimizing the cervical cancer early detection program for the 2015-2019 period. The effort taken was an early detection movement through the method of simultaneous VIA examination in all parts of Indonesia on April 21, 2015. This movement will last for five years and it is expected that by 2019 the number of women of childbearing age who are carried out early detection reaches 50 percent [2].

Mothers' awareness and preparedness with respect to the prevention of cervical cancer in their adolescent daughters were low and inadequate. Mothers should be informed and motivated to play a role in the education of their daughters regarding cervical cancer prevention. Strategies for disseminating information regarding early cervical cancer prevention for adolescent girls are recommended by communicating with both the girls and their mothers and providing them with education regarding cervical cancer prevention [7]. This study aimed to determine the factors that influence mother in behavior of cervical cancer prevention.

14

2. RESEARCH METHOD

This study used analytic survey with cross-sectional design. This study conducted on housewives in the Kota Yogyakarta and Sleman who were members of the Family Empowerment and Welfare amounting to 350 people. This study was conducted in 2016. The population target in this study was housewives who were members of the Family Empowerment and Welfare to manage the confounding variables. To determine the sample size, we used the formula for sample size with calculation of proportion as per Lemeshow with minimum sample size obtained was 350 respondent. Proportional sampling was used to determine the sample chosen. Variables in this study used the theoretical framework approach of Theory of Precede-Proceed (Lawrence Green) [8]. Independent variable consist of the characteristic of respondent (age, parity, education, family income), knowledge level about VIA, and mother' attitude toward VIA. The dependent variable was cervical cancer prevention behavior in mother. This study took primary data. Instrument used in this study was questionnaire. Questionnaire used was development from questionnaire about mothers' knowledge, attitude and behavior toward VIA which have been tested the validity and reliability at Bantul district.

3. RESULTS AND DISCUSSION

This study conducted on 350 respondents was housewives were members of the Family Empowerment and Welfare. The characteristic of the respondent showed in Table 1. Table 1 shows that the major proportion of respondents were aged >35 years old (76.6%), with the highest parity was secundipara in the amount of 166 respondents (47.5%). More than half of the respondents with secondary education amounted to 191 respondents (54.6%). Family income >IDR 1,450,000,00 as many as 189 respondents (54%).

Table 1. Respondent's Characteristic

Characteristic	Amount (n=350)	Percentage (%)
Age		
≤35 years	82	23.4
>35 years	268	76.6
Parity		
Nulipara	6	1.7
Primipara	83	23.7
Secundipara	166	47.5
Multi gravida	95	27.1
Education		
Basic	91	26.0
Medium	191	54.6
High	68	19.4
Family Income		
> IDR 1,450,000,00	189	54.0
≤IDR 1,450,000,00	161	46.0

Table 2 shows that most respondent have not enough knowledge about VIA (80.6%). However, there are still 35 respondents (10%) with a lack of knowledge. The role of health workers is needed in providing insight to mothers regarding early detection of cervical cancer (VIA). Most respondent have

positive attitude toward VIA (54.6%). The behavior of the VIA inspection there are 192 respondents (54.9%) for not VIA.

Table 2. Mothers' Knowledge Level, Attitude, and Behavior toward VIA Test

Characteristic	Amount (n=350)	Percentage (%)
Knowledge level about VIA		
Good	33	9.4
Adequate	282	80.6
Poor	35	10.0
Mothers' attitude toward VIA		
Positive	191	54.6
Negative	159	45.4
Mothers' behavior toward VIA		
Do VIA	158	45.1
Not Do VIA	192	54.9

Bivariate analysis was used to analyze independent and dependent variables. The relation between independent and dependent variables is showed in Table 3. Table 3 shows that factors relating to the behavior of VIA with a p-value <0.05 was educational p=0.024, p=0.036 level of knowledge, and attitudes toward VIA inspection p=0.002. 268 respondents with age>35 years as many as 147 respondents (54.9%) did not do VIA. 166 respondents with secundipara parity as many as 90 respondents (54.2%) did not do VIA. 189 respondents with family income >Rp 1.450.000,00 as many as 110 respondents (58.2%) did not do an VIA. So it can be concluded that there is no significant relationship between parity and family income with IVA behavior (p-value 0.816 and 0.174).

Table 3. The Relationship between Mothers' Characteristic, Knowledge Level, Attitude, and Behavior toward VIA

Variable	Behavior toward VIA				Total		p-value
	Not Do VIA n=192		Do VIA n=158		n=350		
	n	%	n	%	n	%	
Age							
≤35 years	45	54.9	37	45.1	82	100	
> 35 years	147	54.9	121	45.1	268	100	
Parity							
Nulipara	3	50	3	50	6	100	0.816
Primipara	48	57.8	35	42.2	83	100	
Sekundipara	90	54.2	76	45.8	166	100	
Multipara	51	53.7	44	46.3	95	100	
Education Level							
Basic	57	62.6	34	37.4	91	100	0.024*
Medium	102	53.4	89	46.6	191	100	
High	33	48.5	35	51.5	68	100	
Family Income							
> Rp.1.450.000,00	110	58.2	79	41.8	189	100	0.174
≤Rp.1.450.000,00	82	51	79	49	161	100	
Knowledge level							
Superior	16	48.5	17	51.5	33	100	0.036*
Satisfactory	150	53.2	132	46.8	282	100	
Unsatisfactory	26	74.3	9	25.7	35	100	
Attitude toward VIA							
Positive	119	62.3	72	37.7	191	100	0.002*
Negative	73	45.9	86	54.1	159	100	

Multivariate analysis were done to independent variables that have p <0.002. The result of multivariate analysis showed in Table 4. Table 4 shows that the results of statistical tests with logistic regression found that the attitude has a p-value of 0.002. This shows that the factor that most influences the behavior toward VIA is the attitude of respondents.

Cervical cancer could be prevented by screening VIA. Previous research showed that women between the age group 31 to 40 years had more awareness about cervical cancer [9]. Respondents are older than 30 years have a possibility of having a precancerous lesion is higher than under 30 years of age so that respondents in that age will find it more important to conduct tests VIA [10]. The results of this study found

no correlation of age to the prevention behavior of IVA cervical cancer ($p=0$). The results of this study are in accordance with Gustiana D's study that there was no significant relationship between age and early detection of cervical cancer $p=0.306$. It can be associated with susceptibility to disease. Age can not be used as a benchmark for a person to prevent cervical cancer. This can be due to ignorance, no complaints, nor do to prevent cervical cancer has not been necessary [11].

Table 4. Multivariate Analysis Result

	B	Sig.	Exp(B)	95.0% C.I. for EXP(B)	
				Lower	Upper
Attitude toward VIA test	-.666	.002	.514	.335	.788
Constant	.164	.303	1.178		

The results of previous studies conducted by Yuliwati also showed that there was no significant relationship between age and maternal behavior to check VIA [12]. However, different research results Eva who showed that age has a significant relationship with the knowledge that motivates behavior of women for the early detection of cervical cancer by VIA in the district of Central Bogor $p=0.001$ [13].

The results of Heni P et al's research showed that parity had a significant relationship to precancerous cervical lesions [14]. While the chi square test results in this study indicate that there is no significant relationship between parity against the IVA behavior $p=0.816$. The results of this study are inconsistent with Dinengsih research and Sitanggang stated that there is a significant relationship between parity with IVA inspection. A third parity or more are at risk of developing cervical cancer. Multiparity suspected of causing a decrease in endurance. So it is necessary to do the IVA method to detect early cervical cancer [15].

The level of education a person can support or influence a person's level of knowledge and a low level of education is always related to information and knowledge is limited, the higher one's education the higher one's own understanding and knowledge of information obtained would be even higher [16]. Based on the results of this study indicate that there is a significant relationship between education with VIA behavior $p=0.024$. This research is in line with research conducted by Jia on several women in the city of Wufeng, China who explained that education affects the implementation of early detection of cervical cancer by 0.00 ($p=0.05$) with the most characteristics at the low level of education (48.4%) who are willing to carry out early detection of cervical cancer [17]. This is also confirmed by the results of further research by Eva, explains that the education factor has a significant relationship with the level of knowledge possessed by women of childbearing age, where the group of higher education have a proportion of knowledge both larger than the group of secondary education and low [13]. The level of education is one of the factors that determine a person's knowledge and perception of the importance of a case, including the importance of early detection of cervical cancer, caused by highly educated person will be broader view and more receptive to the ideas and new ways of life. It can be concluded that highly educated people will make early detection of cervical cancer [18].

The results showed that no significant relationship between family income on behavior IVA $p=0.174$. The results are consistent with research Gustiana states that there is no relationship to the economic status of cervical cancer prevention behavior ($p=0.561$), research found that respondents with lower economic status also have a good preventive health behaviors [11]. Another study conducted by Wahyuni states that there is no economic status relationship to the behavior of early detection of cervical cancer [19]. Socio-economic conditions affecting the health status change process as it affects the thoughts or beliefs that can lead to changes in health behavior. The incidence of cervical cancer is twice as large in women who have low social classes. In contrast to the results of research Febriani, women who have high incomes likely better prevention of cervical cancer, compared to women with low incomes [17].

Increased knowledge can change people's behavior from negative to positive, in addition to the knowledge also build trust [11]. Having knowledge of a disease, the concept of a disease will be formed in the individual so that it will determine a person's health behavior [20]. Lack of related knowledge about cervical cancer can be an important factor for the high incidence of cervical cancer [9]. Based on the results of chi-square test showed that there is a relationship with the level of knowledge of the behavior of IVA $p=0.036$. The results are consistent with research conducted by Yao Jia on women in the town of Wufeng China stated there is a significant relationship between knowledge with behavioral examination early detection of cervical cancer through IVA method with $p=0.000$ [17]. In addition, research conducted by Nesrin, Turkish women gained knowledge influence the early detection of cervical cancer ($p=0.001$) [21]. Based on research conducted by Khosidah, good knowledge possessed by women of childbearing age about

cervical cancer and its examination can be a motivating factor to try to avoid cervical cancer. Various sources of information that can now be easily accessed by women of childbearing age allow knowledge of cervical cancer and the examination will be better. Lack of knowledge and awareness of the importance of examination is an inhibiting factor for screening for cervical cancer [22].

Attitude is one of the personality elements that a person has to determine his actions and behave towards an object accompanied by positive and negative feelings. Attitudes influence the formation of interest because of the tendency in the subject to accept or reject an object that is good or not [23]. The chi square test results show that there is a significant relationship between attitudes toward behavior IVA $p=0.002$. The results of this study are in line with Febriani's research the attitude of adult women in having a significant relationship with the examination of early detection of cervical cancer $p=0.025$ [18]. TRA explains that beliefs can affect attitudes and social norms which would change the form of the desire well behaved guided or happen in an individual's behavior. This theory confirms the role of one's intention in determining whether a behavior will occur [24].

4. CONCLUSION

Respondent joined this study were mostly aged >35 years old, secundipara, medium level of knowledge, family income more than minimum income regional. There is a relationship of education, level of knowledge, and attitudes with cervical cancer prevention behavior. There is no correlation between age, parity, and family income with the mother's behavior on VIA tests. Attitudes towards VIA tests is the most affect to behavior of the VIA.

ACKNOWLEDGEMENTS

We would like to thank the women and research assistants who participated in this study. The study was funded by Health Polytechnic Ministry of Health Yogyakarta, Indonesia.

REFERENCES

- [1] World Health Organization (WHO). Care of the Preterm and Low-Birth-Weight Newborn World Prematurity Day "Let Them Thrive" 1-5, 2017.
- [2] Septarini NW, Kurniati DPY, Wiryanthini IAD, Putra AWGAE, Sutarga LM. "Initiation of Reproductive Health Care Village Cadres and the Early Detection of Cervical Cancer in the Village of Bangli Bali District- Inisiasi Kader Desa Peduli Kesehatan Reproduksi (KDPKR) Serta Deteksi Dini Kanker Leher Rahim Didesa Pengotan Kabupaten Bangli Bali," *Bul Udayana Mengabdi*. 16(September):107-115, 2017.
- [3] Ministry of Health Republik Indonesia, Cancer Situation-Situasi Penyakit Kanker, Data and Information Center-InfoDATIN, 6-9, 2015.
- [4] Nordianti ME, Wahyono B. "Determinants of Visual Acetate Acid Inspection Visits in Semarang City Health Center-Determinan Kunjungan Inspeksi Visual Asam Asetat di Puskesmas Kota Semarang" *HIGEIA*, 2(1):33-44, 2018.
- [5] Gustafsson L, Ponten J, Bergstrom R, Adami HO. "International Incidence Rates of Invasive Cervical Cancer before Cytological Screening" *International Journal Cancer*. 71:159-165, 1997.
- [6] Gustafsson L, Ponten J, Zack M, Adami HO. "International Incidence Rates of Invasive Cervical Cancer After Introduction of Cytological Screening" *Cancer Causes Control*. 8:755-763, 1997.
- [7] Gustafsson L, Ponten J, Bergstrom R, Adami HO. "International Incidence Rates of Invasive Cervical Cancer Before Cytological Screening" *International Journal Cancer*. 71:159-165, 1997.
- [8] Green LW. *Health Promotion Planning an Educational And Environmental Approach*. Mountain View Toronto - London: Mayfield Publishing Company; 2000.
- [9] Kim HW, Kim DH. "Awareness of Cervical Cancer Prevention among Mothers of Adolescent Daughters in Korea: Qualitative Research" *BMJ Open*. 2015;1-10.
- [10] Abudukadeer A, Azam S, Mutailipu AZ, Qun L, Guilin G, Mijiti S. "Knowledge and Attitude of Uyghur Women in Xinjiang Province of China Related to the Prevention and Early Detection of Cervical Cancer" *World J Surg Oncol*. 13(110):1-7, 2015.
- [11] Mulyati S, Suwarsa O, D A IF. "Effect of Film Media on Mother's Attitude on Early Detection of Cervical Cancer- Pengaruh Media Film terhadap Sikap Ibu pada Deteksi Dini Kanker Serviks" *Jurnal Kesehatan Masyarakat*. 2015;11(2):16-24.
- [12] Gustiana D, Dewi YI, Nurchayati S, "Factors Associated with the Behavior of Prevention of Cervical Cancer in Women of Childbearing Age-Faktor-Faktor yang Berhubungan dengan Perilaku Pencegahan Kanker Serviks Pada Wanita Usia Subur" *JOM PSIK*. 2014;1(2):1-8.
- [13] Yuliwati. "Factors Associated with the Behavior of Wus in Early Detection of Leher Cancer Uterus IVA Method in Prembun Health Center 2012-Faktor-Faktor Yang Berhubungan Dengan Perilaku Wus Dalam Deteksi Dini Kanker

- Lehar Rahim Metode IVA Di Wilayah Puskesmas Prembun 2012” 2012; Available from: <http://repository.ui.ac.id>.
- [14] Sulistiawati E. “Knowledge of Risk Factors, Behavior and Early Detection of Cervical Cancer with IVA in Women in Bogor Tengah District, Bogor-Pengetahuan tentang Faktor Risiko, Perilaku Dan Deteksi Dini Kanker Serviks dengan Inspeksi Visual Asam Asetat (IVA) pada Wanita di Kecamatan Bogor Tengah, Bogor” *Bul Penelit Kesehat.* 42(3):192–202, 2014.
- [15] Purwaningsih H, Prajatmo H. “Risk Factors Associated with the Occurrence of Precancerous Cervical Lesions at Health Centres of Karanganyar Regency” *International Journal Community Medical and Public Health.* 3(6):1437–41, 2016.
- [16] Dinengsih S, Sitanggang E. “Analysis of Early Behavior Detection of Cervical Cancer by IVA Method-Analysis Faktor Perilaku Deteksi Dini Kanker Serviks dengan Metode IVA (Inspeksi Visual Asam Asetat)” *URECOL.* 37–45, 2018.
- [17] Parapat FT, S H S, Sc M, Saraswati LD, Epid M. “Factors Associated with Behavior of Early Detection of Cervical Cancer Visual Acetate Inspection Methods in Candirotto Health Center Temanggung Regency-Faktor Faktor yang Berhubungan dengan Perilaku Deteksi Dini Kanker Leher Rahim Metode Inspeksi Visual Asam Asetat di Puskesmas Candirotto Kabupaten Temanggung” *Jurnal Kesehatan Masyarakat.* 4(4):363–70, 2016.
- [18] Yao J. “Knowledge about Cervical Cancer and Barriers of Screening Program among Women in Wufeng County, A High-Incidence Region of Cervical Cancer in China” 2013.
- [19] Febriani CA. “Factors Associated with Early Detection of Cervical Cancer in Gisting Subdistrict, Tanggamus Regency, Lampung-Faktor-Faktor yang Berhubungan dengan Deteksi Dini Kanker Leher Rahim di Kecamatan Gisting Kabupaten Tanggamus Lampung” *Jurnal Kesehatan.* 2015;VII(2):228–37, 2015.
- [20] Wahyuni S. “Factors Affecting Early Detection Behavior of Cervical Cancer in Ngampel District, Kendal Regency, Central Java-Faktor-Faktor yang Mempengaruhi Perilaku Deteksi Dini Kanker Serviks di Kecamatan Ngampel Kabupaten Kendal Jawa Tengah”, 2013 Available from: <https://bem.unimus.ac.id>.
- [21] Situmorang. “Relationship between Knowledge and Attitudes with Early Detection Behavior in Patients with Cervical Cancer in RSUP Dr. Kariadi Semarang in 2015-Hubungan Pengetahuan Dan Sikap Dengan Perilaku Deteksi Dini Pada Penderita Kanker Serviks Di RSUP Dr. Kariadi Semarang Tahun 2015.” *Jurnal Kesehatan Masyarakat.* 4(1):76–82, 2016.
- [22] Reis N. “Knowledge, Behavior, and Beliefs for Cervical Cancer and Screening in Turkey.” 2012; Available from: [doi:http://dx.doi.org/10.7314/APJCP.2012.13.4.1463](http://dx.doi.org/10.7314/APJCP.2012.13.4.1463)
- [23] Khosidah A. “Factors Affecting Housewives in Conducting IVA Tests as an Effort to Early Detection of Cervical Cancer-Faktor-Faktor Yang Mempengaruhi Ibu Rumah Tangga Dalam Melakukan Tes IVA Sebagai Upaya Deteksi Dini Kanker Serviks,” *Jurnal Ilmu Kebidanan.* 2015;6(2):94–105.
- [24] Sholihah A. “Relationship between the Attitude of Prevention of Cervical Cancer and Interest in Early Detection Using IVA at Fertile Women (WUS) in Mojolaban Village Sukoharjo-Hubungan Antara Sikap Pencegahan Kanker Serviks Dengan Minat Deteksi Dini Menggunakan IVA Pada WUS Di Desa Mojolaban Sukoharjo,” *Artikel Ilmiah.* 20(1), 2015.

ORIGINALITY REPORT

11%

SIMILARITY INDEX

6%

INTERNET SOURCES

4%

PUBLICATIONS

3%

STUDENT PAPERS

PRIMARY SOURCES

1 Lidiya Gutema Lemu, Biruktawit Fekade Woldu, Natnael Eshetu Teke, Nardos Delelegn Bogale, Ermias Ayalew Wondimenew. "Precancerous Cervical Lesions Among HIV-Infected Women Attending HIV Care and Treatment Clinics in Southwest Ethiopia: A Cross-Sectional Study", International Journal of Women's Health, 2021
Publication 1%

2 umexpert.um.edu.my
Internet Source 1%

3 www.sciepub.com
Internet Source 1%

4 Indrabayu, Andi Rezky Fatmasari, Ingrid Nurtanio. "A Colour Space Based Detection for Cervical Cancer Using Fuzzy C-Means Clustering", Proceedings of the 6th International Conference on Bioinformatics and Biomedical Science, 2017
Publication 1%

5	Student Paper	1 %
6	ocs.unud.ac.id Internet Source	<1 %
7	Harshita Prabhakaran, Judie Arulappan. "Effectiveness of Nurse led structured teaching programme on knowledge and practice of postnatal mothers on low birth weight care", Journal of Neonatal Nursing, 2021 Publication	<1 %
8	ejournal.lucp.net Internet Source	<1 %
9	coxsbazarworld.com Internet Source	<1 %
10	Submitted to Universitas Airlangga Student Paper	<1 %
11	www.cervicalopticalimaging.com Internet Source	<1 %
12	F Mutia, E Elviana, A N Yunisya. "Kalanganyar River, Sidoarjo Regency, Indonesia as a natural urban catalyst, living place or society pest", IOP Conference Series: Earth and Environmental Science, 2020 Publication	<1 %

13

Internet Source

<1 %

14

Submitted to Badan PPSDM Kesehatan
Kementerian Kesehatan

Student Paper

<1 %

15

repo.unikadelasalle.ac.id

Internet Source

<1 %

16

www.cambridge.org

Internet Source

<1 %

17

Ahmed Ibrahim, Arja R Aro, Rask, Eero
Pukkala. "Cervical cancer screening in primary
health care setting in Sudan: a comparative
study of visual inspection with acetic acid and
Pap smear", International Journal of Women's
Health, 2012

Publication

<1 %

18

docplayer.net

Internet Source

<1 %

19

Submitted to Syiah Kuala University

Student Paper

<1 %

20

Submitted to Universitas Riau

Student Paper

<1 %

21

eprints.um.edu.my

Internet Source

<1 %

22

Mukhlisiana Ahmad. "Persepsi tentang Kanker Serviks, Promosi Kesehatan, Motivasi Sehat terhadap Perilaku Pencegahan Kanker Serviks pada Bidan di Wilayah Depok", Jurnal Ilmiah Kesehatan, 2017

Publication

<1 %

23

www.slideshare.net

Internet Source

<1 %

24

Chioma Christie Asuzu, Julian Unegbu, Elizabeth Akin-odanye. "Knowledge, attitude and behaviour of the University of Ibadan women towards cancer of the cervix and its prevention", Psycho-Oncology, 2012

Publication

<1 %

25

Fatimah Dewi Anggraeni, Klesia Benedikta. "GAMBARAN DUKUNGAN SUAMI TERHADAP DETEKSI DINI KANKER SERVIKS PADA PASANGAN USIA SUBUR (PUS) DI DUSUN NGASEM DESA TIMBULHARJO SEWON BANTUL TAHUN 2016", Media Ilmu Kesehatan, 2016

Publication

<1 %

26

Febri Hardiyanti, Johan Harlan, Ema Hermawati. "The Association Between Knowledge and Preventive Behavior of Cervical Cancer Among Woman Employees in The Companies in Jakarta", Indonesian Journal of Cancer, 2020

<1 %

Publication

27

info-pendidikan-edukasi.blogspot.com

Internet Source

<1 %

28

nfnc.org.zm

Internet Source

<1 %

29

tampub.uta.fi

Internet Source

<1 %

30

www.koreascience.kr

Internet Source

<1 %

Exclude quotes Off

Exclude matches Off

Exclude bibliography Off