

CHARACTERISTICS OF STUNTING IN TODDLERS AGED 12-59 MONTHS IN KULON PROGO, YOGYAKARTA, INDONESIA

by Yuliasti Eka Purnamaningrum

Submission date: 25-May-2023 08:40PM (UTC+1000)

Submission ID: 2101545058

File name: B51_CH_1-1.pdf (1.03M)

Word count: 348

Character count: 2070

ISSN 2580-9296 (ONLINE)



www.icash.inschool.id



Abstracts of International Conference on Applied Science and Health No. 04, 2019

“Science for mankind: Translating research results into policy and practices”

Faculty of Graduate Studies Mahidol University
Nakhon Pathom, 23rd - 24th July 2019



International Conference on Applied Science and Health
No. 04, 2019

ICASH-C015

CHARACTERISTICS OF STUNTING IN TODDLERS AGED 12-59 MONTHS IN KULON PROGO, YOGYAKARTA, INDONESIA

Yuliasti Eka Purnamaningrum^{1*}, Margono¹, Regita Cita Puspitasari^{1*}, Riska Rahmawati¹,
Risma Fitria Dianasari¹, Izzudien Sobri²

¹⁾ Midwifery Department Health Polytechnic Ministry of Health in Yogyakarta

²⁾ Nutrition Department Health Polytechnic Ministry of Health in Yogyakarta

*Corresponding author's email: yuliasti.eka.purnamaningrum@gmail.com (YEP),
regitacita26@gmail.com (RCP)

ABSTRACT

Introduction: Stunting is a health problem that is caused due to not optimal health conditions and nutritional intake of children during the growth process. Kulon Progo is one of the districts in DIY that is registered in the program of 100 priority districts for stunting intervention. This study aims to determine the characteristic of the stunted toddlers aged 12-59 months in Kulon Progo, Yogyakarta, Indonesia

Method: This study used a cross sectional design. This study involved 84 families with stunting children aged 12-59 months. Data was obtained from interviews and questionnaires, and were then analyzed using the descriptive analysis.

Results: The results show that the stunted children mostly live with families who smoke (61.90%), were born with gestational age > 37 weeks (85.71%) and weight less than 2500 gr (86, 90%). The majority of the stunted toddlers were male (52.38%), and weaned at less than 24 months (53.57%).

Conclusion: This study presents the characteristics of stunted toddlers aged 12-59 months in Kulon Progo, Yogyakarta, Indonesia. The results of this study are expected to be an important input to provide further intervention in improving the nutritional intake for the stunted children.

Keywords: Stunting, Toddlers, Characteristics, Yogyakarta.

CHARACTERISTICS OF STUNTING IN TODDLERS AGED 12-59 MONTHS IN KULON PROGO, YOGYAKARTA, INDONESIA

ORIGINALITY REPORT

13%

SIMILARITY INDEX

0%

INTERNET SOURCES

13%

PUBLICATIONS

0%

STUDENT PAPERS

MATCH ALL SOURCES (ONLY SELECTED SOURCE PRINTED)

6%

★ Karningsih Sudiro, Sri Mulyati. "Baby Spa Effect on Growth", Asian Journal of Applied Sciences, 2018

Publication

Exclude quotes On

Exclude matches Off

Exclude bibliography On

CHARACTERISTICS OF STUNTING IN TODDLERS AGED 12-59 MONTHS IN KULON PROGO, YOGYAKARTA, INDONESIA

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2
