

## Hasil SPSS

### Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
Jenis kelamin * terjadinyastunting	152	100.0%	0	.0%	152	100.0%
Riwayat BBLR * terjadinyastunting	152	100.0%	0	.0%	152	100.0%
Riwayat ASI Eksklusif * terjadinyastunting	152	100.0%	0	.0%	152	100.0%
Riwayat KEK * terjadinyastunting	152	100.0%	0	.0%	152	100.0%
Riwayat ANC * terjadinyastunting	152	100.0%	0	.0%	152	100.0%

## Jenis Kelamin

### Crosstabulation

		Terjadinya Stunting		Total
		Kasus	Kelompok kontrol	
Jenis kelamin	Perempuan	Count 24	Count 34	Count 58
	% within jenis kelamin	15.7%	22.3%	38.1%
Laki-laki	Count	52	42	94
	% within jenis kelamin	34.3%	7.7%	61.9%
Total	Count	76	76	152
	% within jenis kelamin	50.0%	50.0%	100.0%

### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	.046 <sup>a</sup>	1	.004		
Continuity Correction <sup>b</sup>	2.750	1	.009		
Likelihood Ratio	1.782	1	.005		
Fisher's Exact Test				.007	.006
Linear-by-Linear Association	2.278	1	.004		
N of Valid Cases <sup>b</sup>	150				

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 5.00.

b. Computed only for a 2x2 table

### Symmetric Measures

		Value	Approx. Sig.
Nominal by Nominal	Contingency Coefficient	.229	.004
N of Valid Cases		152	

### Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for jenis kelamin (perempuan = laki-laki)	0.679	0.894	1.214
For cohort terjadinya stunting	.046	.798	4.926
For cohort terjadinya stunting	.444	.286	.692
N of Valid Cases	152		

### Riwayat BBLR

#### Crosstabulation

			Terjadinya Stunting		Total
			Kasus	Kelompok kontrol	
Riwayat BBLR	Beresiko	Count	16	13	29
		% within riwayat BBLR	10.5%	8.6%	19.1%
	Tidak beresiko	Count	60	63	123
		% within riwayat BBLR	39.5%	41.4%	80.9%
Total		Count	76	76	152
		% within riwayat BBLR	50.0%	50.0%	100.0%

#### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	1.432 <sup>a</sup>	1	.004		
Continuity Correction <sup>b</sup>	6.217	1	.013		
Likelihood Ratio	7.280	1	.007		
Fisher's Exact Test				.009	.006
Linear-by-Linear Association	7.068	1	.008		
N of Valid Cases <sup>b</sup>	152				

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 5.00.

b. Computed only for a 2x2 table

#### Symmetric Measures

		Value	Approx. Sig.
Nominal by Nominal	Contingency Coefficient	.299	.004
N of Valid Cases		152	

### Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for Riwayat BBLR (beresiko = tidak beresiko)	.004	1.691	6.906
For cohort terjadinya stunting	.001	1.432	1.714
For cohort terjadinya stunting	.117	.062	.221
N of Valid Cases	152		

### Riwayat ASI Eksklusif

#### Crosstabulation

		Terjadinya Stunting		Total
		Kasus	Kelompok kontrol	
Riwayat ASI Beresiko eksklusif	Count	25	19	44
	% within riwayat ASI Eksklusif	16.4%	12.5%	29.0%
Tidak beresiko	Count	51	57	108
	% within riwayat ASI Eksklusif	33.6%	37.5%	71.0%
Total	Count	76	76	152
	% within riwayat ASI Eksklusif	50,0%	50,0%	100,0%

#### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	1.169 <sup>a</sup>	1	.003		
Continuity Correction <sup>b</sup>	6.565	1	.000		
Likelihood Ratio	8.850	1	.000		
Fisher's Exact Test				.000	.000
Linear-by-Linear Association	8.836	1	.000		
N of Valid Cases <sup>b</sup>	152				

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 5.00.

b. Computed only for a 2x2 table

#### Symmetric Measures

		Value	Approx. Sig.
Nominal by Nominal	Contingency Coefficient	.497	.003
N of Valid Cases		152	

### Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for Riwayat ASI Eksklusif (beresiko = tidak beresiko)	.003	1.027	1,547
For cohort terjadinya stunting	.002	.992	2.272
For cohort terjadinya stunting	.208	.128	.337
N of Valid Cases	152		

### Riwayat KEK selama Hamil

#### Crosstabulation

			Terjadinya Stunting		Total
			Kasus	Kelompok kontrol	
Riwayat KEK	Beresiko	Count	39	19	58
		% within riwayat KEK	25.7%	15.2%	38.1%
	Tidak beresiko	Count	37	57	94
		% within riwayat KEK	24.3%	34.8%	61.9%
Total		Count	76	76	152
		% within riwayat KEK	50.0%	50.0%	100.0%

#### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	1.583 <sup>a</sup>	1	.000		
Continuity Correction <sup>b</sup>	5.085	1	.000		
Likelihood Ratio	6.266	1	.000		
Fisher's Exact Test				.000	.000
Linear-by-Linear Association	6.523 <sup>a</sup>	1	.000		
N of Valid Cases <sup>b</sup>	152				

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 5.00.

b. Computed only for a 2x2 table

#### Symmetric Measures

		Value	Approx. Sig.
Nominal by Nominal	Contingency Coefficient	.632	.000
N of Valid Cases		152	

### Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for Riwayat KEK (beresiko = tidak beresiko)	.000	1.067	1.765
For cohort terjadinya stunting	.002	.348	2.719
For cohort terjadinya stunting	.595	.362	.979
N of Valid Cases	152		

### Riwayat ANC Ibu Selama Hamil

#### Crosstabulation

			Terjadinya Stunting		Total
			Kasus	Kelompok kontrol	
Riwayat ANC	Beresiko	Count	21	16	37
		% within riwayat ANC	13.8%	10.5%	24.3%
	Tidak beresiko	Count	55	60	115
		% within riwayat ANC	36.2%	39.5%	75.7%
Total		Count	76	76	152
		% within riwayat ANC	50.0%	50.0%	100.0%

#### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	1.282 <sup>a</sup>	1	.001		
Continuity Correction <sup>b</sup>	17.826	1	.001		
Likelihood Ratio	16.478	1	.000		
Fisher's Exact Test				.001	.000
Linear-by-Linear Association	20.827	1	.000		
N of Valid Cases <sup>b</sup>	152				

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 5.00.

b. Computed only for a 2x2 table

#### Symmetric Measures

		Value	Approx. Sig.
Nominal by Nominal	Contingency Coefficient	.407	.000
N of Valid Cases		152	

**Risk Estimate**

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for Riwayat ANC (beresiko = tidak beresiko)	.001	1.205	1.459
For cohort terjadinya stunting	.003	.557	4.797
For cohort terjadinya stunting	.146	.059	.360
N of Valid Cases	152		