

# **LAMPIRAN**

## Lampiran 1

## ANGGARAN PENELITIAN

No	Kegiatan	Volume	Satuan	Unit cost	Jumlah
1.	Biaya penelitian				
	a. Biaya studi pendahuluan	5	Kali	100.000	500.000
	b. Biaya pengambilan data	200	RM	1000	102.500
2.	Transportasi penelitian Transportasi ke lokasi	22	Kali	10.000	220.000
3.	ATK dan pengadaan (pencetakan dan penjiilidan)				
	a. Penyusunan skripsi	4	Paket	20.000	80.000
	b. Seminar				
	c. Revisi	5	Paket	20.000	100.000
	d. Pengajuan etik penelitian	4	Paket	20.000	80.000
		1	Paket	50.000	50.000
	e. Penyusunan laporan hasil	4	Paket	25.000	120.000
	f. Sidang hasil skripsi				
	g. Revisi laporan skripsi akhir	5	Paket	25.000	125.000
		4	Paket	25.000	120.000
	<b>Jumlah</b>				<b>1.497.500</b>

Lampiran 2

**JADWAL PENELITIAN**

NO	KEGIATAN	BULAN																											
		NOVEMBER				DESEMBER				JANUARI				FEBRUARI				MARET				APRIL				MEI			
		1	2	3	4	1	2	3	4	1	2	3	4	1	2	3	4	1	2	3	4	1	2	3	4	1	2	3	4
1	Penyusunan Proposal Skripsi	■																											
2	Seminar Proposal Skripsi									■																			
3	Revisi Proposal Skripsi										■																		
4	Perijinan Penelitian											■	■	■	■	■	■												
5	Persiapan Penelitian														■	■	■												
6	Pelaksanaan Penelitian																	■	■	■	■	■	■	■	■				
7	Pengolahan Data																											■	
8	Laporan Skripsi																										■	■	■
9	Sidang Skripsi																												■
10	Revisi Laporan Skripsi																												■





Lampiran 5

**A. HASIL ANALISIS UNIVARIAT**

**Frequencies**

**Statistics**

Kejadian Preeklampsia

N	Valid	194
	Missing	0

**Kejadian Preeklampsia**

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid Preeklampsia	97	50,0	50,0	50,0
Tidak preeklampsia	97	50,0	50,0	100,0
Total	194	100,0	100,0	

**Frequencies**

**Statistics**

Usia Ibu

N	Valid	194
	Missing	0

**Usia Ibu**

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid berisiko (usia <20 atau >35 tahun)	48	24,7	24,7	24,7
tidak berisiko (usia 20-35 tahun)	146	75,3	75,3	100,0
Total	194	100,0	100,0	

**Frequencies**

**Statistics**

Paritas

N	Valid	194
	Missing	0

**Paritas**

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid berisiko (>3 kali)	18	9,3	9,3	9,3
tidak berisiko (1-3 kali)	176	90,7	90,7	100,0
Total	194	100,0	100,0	

**Frequencies****Statistics**

Riwayat Preeklampsia Sebelumnya

N	Valid	194
	Missing	0

**Riwayat Preeklampsia Sebelumnya**

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Ada	17	8,8	8,8	8,8
	Tidak ada	177	91,2	91,2	100,0
	Total	194	100,0	100,0	

**Frequencies****Statistics**

Diabetes Melitus

N	Valid	194
	Missing	0

**Diabetes Melitus**

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Ada	12	6,2	6,2	6,2
	Tidak ada	182	93,8	93,8	100,0
	Total	194	100,0	100,0	

**Frequencies****Statistics**

Obesitas

N	Valid	194
	Missing	0

**Obesitas**

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Ada	53	27,3	27,3	27,3
	Tidak ada	141	72,7	72,7	100,0
	Total	194	100,0	100,0	

**Frequencies****Statistics**

Hipertensi Kronik

N	Valid	194
	Missing	0

### Hipertensi Kronik

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid				
Ada	25	12,9	12,9	12,9
Tidak ada	169	87,1	87,1	100,0
Total	194	100,0	100,0	

## B. HASIL ANALISIS BIVARIAT

### Crosstabs

#### Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
Usia Ibu * Kejadian Preeklampsia	194	100,0 %	0	0,0%	194	100,0%

#### Usia Ibu \* Kejadian Preeklampsia Crosstabulation

		Kejadian Preeklampsia		Total
		Preeklampsia	Tdk preeklampsia	
Usia Ibu berisiko (usia <20 atau >35 tahun)	Count	36	12	48
	% within Usia Ibu	75,0%	25,0%	100,0%
tidak berisiko (usia 20-35 tahun)	Count	61	85	146
	% within Usia Ibu	41,8%	58,2%	100,0%
Total	Count	97	97	194
	% within Usia Ibu	50,0%	50,0%	100,0%

### Chi-Square Tests

	Value	df	Asymptotic Significance (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	15,945 <sup>a</sup>	1	,000		
Continuity Correction <sup>b</sup>	14,644	1	,000		
Likelihood Ratio	16,521	1	,000		
Fisher's Exact Test				,000	,000
Linear-by-Linear Association	15,863	1	,000		
N of Valid Cases	194				

a. 0 cells (,0%) have expected count less than 5. The minimum expected count is 24,00.

b. Computed only for a 2x2 table



### Symmetric Measures

		Value	Asymptotic Standard Error <sup>a</sup>	Approximate T <sup>b</sup>	Approximate Significance
Nominal by Nominal	Contingency Coefficient	,276			,000
Interval by Interval	Pearson's R	,287	,066	4,147	,000 <sup>c</sup>
Ordinal by Ordinal	Spearman Correlation	,287	,066	4,147	,000 <sup>c</sup>
N of Valid Cases		194			

a. Not assuming the null hypothesis.

b. Using the asymptotic standard error assuming the null hypothesis.

c. Based on normal approximation.

### Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for Usia Ibu (berisiko (usia <20 atau >35 tahun) / tidak berisiko (usia 20-35 tahun))	4,180	2,012	8,687
For cohort Kejadian Preeklampsia = Preeklampsia	1,795	1,396	2,309
For cohort Kejadian Preeklampsia = Tdk preeklampsia	,429	,258	,714
N of Valid Cases	194		

### Crosstabs

#### Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
Paritas * Kejadian Preeklampsia	194	100,0%	0	0,0%	194	100,0%

#### Paritas \* Kejadian Preeklampsia Crosstabulation

			Kejadian Preeklampsia		Total
			Preeklampsia	Tdk preeklampsia	
Paritas berisiko (>3 kali)	Count		14	4	18
	% within Paritas		77,8%	22,2%	100,0%
tidak berisiko (1-3 kali)	Count		83	93	176
	% within Paritas		47,2%	52,8%	100,0%
Total	Count		97	97	194
	% within Paritas		50,0%	50,0%	100,0%

### Chi-Square Tests

	Value	df	Asymptotic Significance (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	6,124 <sup>a</sup>	1	,013		
Continuity Correction <sup>b</sup>	4,960	1	,026		
Likelihood Ratio	6,452	1	,011		
Fisher's Exact Test				,024	,012
Linear-by-Linear Association	6,092	1	,014		
N of Valid Cases	194				

a. 0 cells (,0%) have expected count less than 5. The minimum expected count is 9,00.

b. Computed only for a 2x2 table

### Symmetric Measures

	Value	Asymptotic Standard Error <sup>a</sup>	Approximate T <sup>b</sup>	Approximate Significance
Nominal by Nominal Contingency Coefficient	,175			,013
Interval by Interval Pearson's R	,178	,063	2,502	,013 <sup>c</sup>
Ordinal by Ordinal Spearman Correlation	,178	,063	2,502	,013 <sup>c</sup>
N of Valid Cases	194			

a. Not assuming the null hypothesis.

b. Using the asymptotic standard error assuming the null hypothesis.

c. Based on normal approximation.

### Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for Paritas (berisiko (>3 kali) / tidak berisiko (1-3 kali))	3,922	1,242	12,385
For cohort Kejadian Preeklampsia = Preeklampsia	1,649	1,231	2,209
For cohort Kejadian Preeklampsia = Tdk preeklampsia	,421	,175	1,009
N of Valid Cases	194		

### Crosstabs

#### Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
Riwayat Preeklampsia Sebelumnya *	194	100,0%	0	0,0%	194	100,0%
Kejadian Preeklampsia						

**Riwayat Preeklampsia Sebelumnya \* Kejadian Preeklampsia Crosstabulation**

			Kejadian Preeklampsia		Total
			Preeklampsi a	Tdk preeklampsia	
Riwayat Preeklampsia Sebelumnya	Ada	Count % within Riwayat Preeklampsia Sebelumnya	16 94,1%	1 5,9%	17 100,0%
	Tidak ada	Count % within Riwayat Preeklampsia Sebelumnya	81 45,8%	96 54,2%	177 100,0%
Total		Count	97	97	194
		% within Riwayat Preeklampsia Sebelumnya	50,0%	50,0%	100,0%

**Chi-Square Tests**

	Value	df	Asymptotic Significance (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	14,506 <sup>a</sup>	1	,000		
Continuity Correction <sup>b</sup>	12,637	1	,000		
Likelihood Ratio	17,233	1	,000		
Fisher's Exact Test				,000	,000
Linear-by-Linear Association	14,432	1	,000		
N of Valid Cases	194				

a. 0 cells (,0%) have expected count less than 5. The minimum expected count is 8,50.

b. Computed only for a 2x2 table

**Symmetric Measures**

		Value	Asymptotic Standard Error <sup>a</sup>	Approximate T <sup>b</sup>	Approximate Significance
Nominal by Nominal	Contingency Coefficient	,264			,000
Interval by Interval	Pearson's R	,273	,048	3,939	,000 <sup>c</sup>
Ordinal by Ordinal	Spearman Correlation	,273	,048	3,939	,000 <sup>c</sup>
N of Valid Cases		194			

a. Not assuming the null hypothesis.

b. Using the asymptotic standard error assuming the null hypothesis.

c. Based on normal approximation.

### Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for Riwayat Preeklampsia Sebelumnya (Ada / Tidak ada)	18,963	2,461	146,101
For cohort Kejadian Preeklampsia = Preeklampsia	2,057	1,684	2,511
For cohort Kejadian Preeklampsia = Tdk preeklampsia	,108	,016	,730
N of Valid Cases	194		

### Crosstabs

#### Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
Diabetes Melitus * Kejadian Preeklampsia	194	100,0%	0	0,0%	194	100,0%

#### Diabetes Melitus \* Kejadian Preeklampsia Crosstabulation

			Kejadian Preeklampsia		Total
			Preeklampsia	Tdk preeklampsia	
Diabetes Melitus	Ada	Count	9	3	12
		% within Diabetes Melitus	75,0%	25,0%	100,0%
	Tidak ada	Count	88	94	182
		% within Diabetes Melitus	48,4%	51,6%	100,0%
Total		Count	97	97	194
		% within Diabetes Melitus	50,0%	50,0%	100,0%

#### Chi-Square Tests

	Value	df	Asymptotic Significance (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	3,198 <sup>a</sup>	1	,074		
Continuity Correction <sup>b</sup>	2,221	1	,136		
Likelihood Ratio	3,337	1	,068		
Fisher's Exact Test				,133	,067
Linear-by-Linear Association	3,181	1	,074		
N of Valid Cases	194				

a. 0 cells (,0%) have expected count less than 5. The minimum expected count is 6,00.

b. Computed only for a 2x2 table

### Symmetric Measures

		Value	Asymptotic Standard Error <sup>a</sup>	Approximate T <sup>b</sup>	Approximate Significance
Nominal by Nominal	Contingency Coefficient	,127			,074
Interval by Interval	Pearson's R	,128	,065	1,794	,074 <sup>c</sup>
Ordinal by Ordinal	Spearman Correlation	,128	,065	1,794	,074 <sup>c</sup>
N of Valid Cases		194			

a. Not assuming the null hypothesis.

b. Using the asymptotic standard error assuming the null hypothesis.

c. Based on normal approximation.

### Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for Diabetes Melitus (Ada / Tidak ada)	3,205	,840	12,221
For cohort Kejadian Preeklampsia = Preeklampsia	1,551	1,083	2,222
For cohort Kejadian Preeklampsia = Tdk preeklampsia	,484	,180	1,303
N of Valid Cases		194	

### Crosstabs

#### Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
Obesitas * Kejadian Preeklampsia	194	100,0%	0	0,0%	194	100,0%

#### Obesitas \* Kejadian Preeklampsia Crosstabulation

			Kejadian Preeklampsia		Total
			Preeklampsia	Tdk preeklampsia	
Obesitas	Ada	Count	42	11	53
		% within Obesitas	79,2%	20,8%	100,0%
	Tidak ada	Count	55	86	141
		% within Obesitas	39,0%	61,0%	100,0%
Total		Count	97	97	194
		% within Obesitas	50,0%	50,0%	100,0%

### Chi-Square Tests

	Value	df	Asymptotic Significance (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	24,948 <sup>a</sup>	1	,000	,000	,000
Continuity Correction <sup>b</sup>	23,364	1	,000		
Likelihood Ratio	26,212	1	,000		
Fisher's Exact Test					
Linear-by-Linear Association	24,819	1	,000		
N of Valid Cases	194				

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 26,50.

b. Computed only for a 2x2 table

### Symmetric Measures

	Value	Asymptotic Standard Error <sup>a</sup>	Approximate T <sup>b</sup>	Approximate Significance
Nominal by Nominal Contingency Coefficient	,338			,000
Interval by Interval Pearson's R	,359	,063	5,323	,000 <sup>c</sup>
Ordinal by Ordinal Spearman Correlation	,359	,063	5,323	,000 <sup>c</sup>
N of Valid Cases	194			

a. Not assuming the null hypothesis.

b. Using the asymptotic standard error assuming the null hypothesis.

c. Based on normal approximation.

### Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for Obesitas (Ada / Tidak ada)	5,970	2,834	12,577
For cohort Kejadian Preeklampsia = Preeklampsia	2,032	1,585	2,604
For cohort Kejadian Preeklampsia = Tdk preeklampsia	,340	,198	,585
N of Valid Cases	194		

### Crosstabs

#### Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
Hipertensi Kronik * Kejadian Preeklampsia	194	100,0%	0	0,0%	194	100,0%

**Hipertensi Kronik \* Kejadian Preeklampsia Crosstabulation**

			Kejadian Preeklampsia		Total
			Preeklampsia	Tdk preeklampsia	
Hipertensi Kronik	Ada	Count % within Hipertensi Kronik	22 88,0%	3 12,0%	25 100,0%
	Tidak ada	Count % within Hipertensi Kronik	75 44,4%	94 55,6%	169 100,0%
Total		Count % within Hipertensi Kronik	97 50,0%	97 50,0%	194 100,0%

**Chi-Square Tests**

	Value	df	Asymptotic Significance (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	16,576 <sup>a</sup>	1	,000	,000	,000
Continuity Correction <sup>b</sup>	14,877	1	,000		
Likelihood Ratio	18,452	1	,000		
Fisher's Exact Test					
Linear-by-Linear Association	16,491	1	,000		
N of Valid Cases	194				

a. 0 cells (,0%) have expected count less than 5. The minimum expected count is 12,50.

b. Computed only for a 2x2 table

**Symmetric Measures**

		Value	Asymptotic Standard Error <sup>a</sup>	Approximate T <sup>b</sup>	Approximate Significance
Nominal by Nominal	Contingency Coefficient	,281			,000
Interval by Interval	Pearson's R	,292	,056	4,235	,000 <sup>c</sup>
Ordinal by Ordinal	Spearman Correlation	,292	,056	4,235	,000 <sup>c</sup>
N of Valid Cases		194			

a. Not assuming the null hypothesis.

b. Using the asymptotic standard error assuming the null hypothesis.

c. Based on normal approximation.

### Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for Hipertensi Kronik (Ada / Tidak ada)	9,191	2,650	31,884
For cohort Kejadian Preeklampsia = Preeklampsia	1,983	1,588	2,477
For cohort Kejadian Preeklampsia = Tdk preeklampsia	,216	,074	,629
N of Valid Cases	194		

## C. HASIL ANALISIS MULTIVARIAT

### Logistic Regression

#### Case Processing Summary

Unweighted Cases <sup>a</sup>		N	Percent
Selected Cases	Included in Analysis	194	100,0
	Missing Cases	0	,0
	Total	194	100,0
Unselected Cases		0	,0
	Total	194	100,0

a. If weight is in effect, see classification table for the total number of cases.

#### Dependent Variable Encoding

Original Value	Internal Value
Preeklampsia	0
Tdk preeklampsia	1

#### Categorical Variables Codings

		Frequency	Parameter coding (1)
Hipertensi Kronik	Ada	25	1,000
	Tidak ada	169	,000
Paritas	berisiko (>3 kali)	18	1,000
	tidak berisiko (1-3 kali)	176	,000
Riwayat Preeklampsia Sebelumnya	Ada	17	1,000
	Tidak ada	177	,000
Diabetes Melitus	Ada	12	1,000
	Tidak ada	182	,000
Obesitas	Ada	53	1,000
	Tidak ada	141	,000
Usia Ibu	berisiko (usia <20 atau >35 tahun)	48	1,000
	tidak berisiko (usia 20-35 tahun)	146	,000



**Block 0: Beginning Block**

**Classification Table<sup>a,b</sup>**

Observed			Predicted		
			Kejadian Preeklampsia		Percentage Correct
			Preeklampsia	Tdk preeklampsia	
Step 0	Kejadian Preeklampsia	Preeklampsia	0	97	,0
	Preeklampsia	Tdk preeklampsia	0	97	100,0
	Overall Percentage				50,0

a. Constant is included in the model.

b. The cut value is ,500

**Variables in the Equation**

	B	S.E.	Wald	df	Sig.	Exp(B)
Step 0 Constant	,000	,144	,000	1	1,000	1,000

**Variables not in the Equation**

			Score	df	Sig.
Step 0	Variables	Usia Ibu	15,945	1	,000
		Riwayat Preeklampsia Sebelumnya	14,506	1	,000
		Diabetes Melitus	3,198	1	,074
		Obesitas	24,948	1	,000
		Hipertensi Kronik	16,576	1	,000
		Paritas	6,124	1	,013
	Overall Statistics		56,567	6	,000

**Block 1: Method = Enter**

**Omnibus Tests of Model Coefficients**

		Chi-square	df	Sig.
Step 1	Step	67,734	6	,000
	Block	67,734	6	,000
	Model	67,734	6	,000

### Model Summary

Step	-2 Log likelihood	Cox & Snell R Square	Nagelkerke R Square
1	201,207 <sup>a</sup>	,295	,393

a. Estimation terminated at iteration number 6 because parameter estimates changed by less than ,001.

### Classification Table<sup>a</sup>

Observed	Predicted		
	Kejadian Preeklampsia		Percentage Correct
	Preeklampsia	Tdk preeklampsia	
Step 1 Kejadian Preeklampsia	Preeklampsia	77	79,4
	Tdk preeklampsia	27	72,2
Overall Percentage			75,8

a. The cut value is ,500

### Variables in the Equation

	B	S.E.	Wald	df	Sig.	Exp(B)	95% C.I.for EXP(B)	
							Lower	Upper
Step 1 <sup>a</sup> Usia Ibu	1,404	,466	9,078	1	,003	4,073	1,634	10,153
Paritas	-,141	,785	,032	1	,858	,869	,187	4,045
Riwayat Preeklampsia Sebelumnya	1,818	1,134	2,569	1	,109	6,157	,667	56,843
Diabetes Melitus	1,691	,746	5,146	1	,023	5,427	1,259	23,399
Obesitas	1,929	,421	20,991	1	,000	6,884	3,016	15,712
Hipertensi Kronik	2,112	,694	9,269	1	,002	8,264	2,122	32,184
Constant	-16,421	3,353	23,990	1	,000	,000		

a. Variable(s) entered on step 1: Usia Ibu, Paritas, Riwayat Preeklampsia Sebelumnya, Diabetes Melitus, Obesitas, Hipertensi Kronik.

### Logistic Regression

#### Case Processing Summary

Unweighted Cases <sup>a</sup>		N	Percent
Selected Cases	Included in Analysis	194	100,0
	Missing Cases	0	,0
	Total	194	100,0
Unselected Cases		0	,0
Total		194	100,0

a. If weight is in effect, see classification table for the total number of cases.

**Dependent Variable Encoding**

Original Value	Internal Value
Preeklampsia	0
Tdk preeklampsia	1

**Categorical Variables Codings**

		Frequency	Parameter coding (1)
Hipertensi Kronik	Ada	25	1,000
	Tidak ada	169	,000
Riwayat Preeklampsia Sebelumnya	Ada	17	1,000
	Tidak ada	177	,000
Diabetes Melitus	Ada	12	1,000
	Tidak ada	182	,000
Obesitas	Ada	53	1,000
	Tidak ada	141	,000
Usia Ibu	berisiko (usia <20 atau >35 tahun)	48	1,000
	tidak berisiko (usia 20- 35 tahun)	146	,000

**Block 0: Beginning Block**

**Classification Table<sup>a,b</sup>**

Observed	Predicted			Percentage Correct
	Kejadian Preeklampsia			
	Preeklampsia	Tdk preeklampsia		
Step 0 Kejadian Preeklampsia	Preeklampsia	0	97	,0
	Tdk preeklampsia	0	97	100,0
Overall Percentage				50,0

a. Constant is included in the model.

b. The cut value is ,500

**Variables in the Equation**

	B	S.E.	Wald	df	Sig.	Exp(B)
Step 0 Constant	,000	,144	,000	1	1,000	1,000

**Variables not in the Equation**

			Score	df	Sig.
Step 0	Variables	Usia Ibu	15,945	1	,000
		Riwayat Preeklampsia Sebelumnya	14,506	1	,000
		Diabetes Melitus	3,198	1	,074
		Obesitas	24,948	1	,000
		Hipertensi Kronik	16,576	1	,000
	Overall Statistics		56,229	5	,000

**Block 1: Method = Enter**

**Omnibus Tests of Model Coefficients**

		Chi-square	df	Sig.
Step 1	Step	67,702	5	,000
	Block	67,702	5	,000
	Model	67,702	5	,000

**Model Summary**

Step	-2 Log likelihood	Cox & Snell R Square	Nagelkerke R Square
1	201,239 <sup>a</sup>	,295	,393

a. Estimation terminated at iteration number 6 because parameter estimates changed by less than ,001.

**Classification Table<sup>a</sup>**

Observed			Predicted		
			Kejadian Preeklampsia		Percentage Correct
			Preeklampsia	Tdk preeklampsia	
Step 1	Kejadian Preeklampsia	Preeklampsia	77	20	79,4
		Tdk preeklampsia	27	70	72,2
	Overall Percentage				75,8

a. The cut value is ,500

### Variables in the Equation

	B	S.E.	Wald	df	Sig.	Exp(B)	95% C.I.for EXP(B)	
							Lower	Upper
Step 1 <sup>a</sup>								
Usia Ibu	1,375	,435	9,974	1	,002	3,955	1,685	9,283
Riwayat Preeklampsia Sebelumnya	1,796	1,128	2,534	1	,111	6,027	,660	55,036
Diabetes Melitus	1,692	,746	5,149	1	,023	5,433	1,259	23,436
Obesitas	1,925	,420	20,964	1	,000	6,856	3,007	15,629
Hipertensi Kronik	2,115	,693	9,310	1	,002	8,293	2,131	32,275
Constant	-16,601	3,212	26,710	1	,000	,000		

a. Variable(s) entered on step 1: Usia Ibu, Riwayat Preeklampsia Sebelumnya, Diabetes Melitus, Obesitas, Hipertensi Kronik.

### Logistic Regression

#### Case Processing Summary

Unweighted Cases <sup>a</sup>		N	Percent
Selected Cases	Included in Analysis	194	100,0
	Missing Cases	0	,0
	Total	194	100,0
Unselected Cases		0	,0
Total		194	100,0

a. If weight is in effect, see classification table for the total number of cases.

#### Dependent Variable Encoding

Original Value	Internal Value
Preeklampsia	0
Tdk preeklampsia	1

#### Categorical Variables Codings

		Frequency	Parameter coding (1)
Hipertensi Kronik	Ada	25	1,000
	Tidak ada	169	,000
Diabetes Melitus	Ada	12	1,000
	Tidak ada	182	,000
Obesitas	Ada	53	1,000
	Tidak ada	141	,000
Usia Ibu	berisiko (usia <20 atau >35 tahun)	48	1,000
	tidak berisiko (usia 20-35 tahun)	146	,000

**Block 0: Beginning Block**

**Classification Table<sup>a,b</sup>**

Observed			Predicted		
			Kejadian Preeklampsia		Percentage Correct
			Preeklampsia	Tdk preeklampsia	
Step 0	Kejadian Preeklampsia	Preeklampsia	0	97	,0
	Preeklampsia Tdk preeklampsia	Tdk preeklampsia	0	97	100,0
Overall Percentage					50,0

a. Constant is included in the model.

b. The cut value is ,500

**Variables in the Equation**

	B	S.E.	Wald	df	Sig.	Exp(B)
Step 0 Constant	,000	,144	,000	1	1,000	1,000

**Variables not in the Equation**

	Score	df	Sig.
Step 0 Variables Usia Ibu	15,945	1	,000
Diabetes Melitus	3,198	1	,074
Obesitas	24,948	1	,000
Hipertensi Kronik	16,576	1	,000
Overall Statistics	54,181	4	,000

**Block 1: Method = Enter**

**Omnibus Tests of Model Coefficients**

	Chi-square	df	Sig.
Step 1 Step	64,200	4	,000
Block	64,200	4	,000
Model	64,200	4	,000

### Model Summary

Step	-2 Log likelihood	Cox & Snell R Square	Nagelkerke R Square
1	204,741 <sup>a</sup>	,282	,376

a. Estimation terminated at iteration number 5 because parameter estimates changed by less than ,001.

### Classification Table<sup>a</sup>

Observed			Predicted		
			Kejadian Preeklampsia		Percentage Correct
			Preeklampsia	Tdk preeklampsia	
Step 1	Kejadian Preeklampsia	Preeklampsia	76	21	78,4
		Tdk preeklampsia	27	70	72,2
	Overall Percentage				75,3

a. The cut value is ,500

### Variables in the Equation

		B	S.E.	Wald	df	Sig.	Exp(B)	95% C.I. for EXP(B)	
								Lower	Upper
Step 1 <sup>a</sup>	Usia Ibu	1,537	,420	13,415	1	,000	4,649	2,043	10,579
	Diabetes Melitus	1,668	,749	4,952	1	,026	5,301	1,220	23,030
	Obesitas	1,995	,414	23,181	1	,000	7,352	3,264	16,560
	Hipertensi Kronik	2,373	,675	12,372	1	,000	10,727	2,859	40,241
	Constant	-13,943	2,482	31,560	1	,000	,000		

a. Variable(s) entered on step 1: Usia Ibu, Diabetes Melitus, Obesitas, Hipertensi Kronik.