

The relationship between fried food consumption and physical activity with diabetes mellitus in Yogyakarta, Indonesia

by Tri Siswati

Submission date: 30-Apr-2021 09:09AM (UTC+0700)

Submission ID: 1573965005

File name: 2017_IJCMPH.pdf (538.15K)

Word count: 4555

Character count: 21490

Original Research Article

DOI: <http://dx.doi.org/10.18203/2394-6040.ijcmph20164709>

The relationship between fried food consumption and physical activity with diabetes mellitus in Yogyakarta, Indonesia

Toto Sudargo^{1*}, Shinta Pertiwi¹, Ruth Audy Alexander¹, Tri Siswati², Yuli Ernawati³

¹Department of Nutrition and Health, Faculty of Medicine, Gadjah Mada University, Yogyakarta, Indonesia

²Health Polytechnic, Ministry of Health, Yogyakarta, Indonesia

³Ners Study Program, School of Health Science Wira Husada, Yogyakarta, Indonesia

Received: 17 October 2016

Accepted: 15 November 2016

*Correspondence:

Dr. Toto Sudargo,

E-mail: toto_sudargo@yahoo.co.id

Copyright: © the author(s), publisher and licensee Medip Academy. This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License, which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ABSTRACT

Background: Diabetes mellitus (DM) is one of degenerative disease and the prevalence is increasing each year. WHO predicted that in 2025 the number of diabetes mellitus outpatients in the world can reach up to 300 millions people. Early detection is needed to prevent higher prevalence type 2 DM. Moreover, nowadays people are tend to have sedentary lifestyle and bad food pattern. The Objective of this research aims to determine risk factors which related the evidence of type 2 DM.

Methods: It was an observational study with cross sectional design. Subjects were 179 people of the study were staffs of Gadjah Mada University who did general check-up at Gadjah Mada Medical Center (GMC) health center on March and April 2014. Research variable included diabetic family history, ethnic, physical activity, habit of smoking history, obese, eating pattern, fried foods consumption and diabetes. Data of eating pattern and fried foods consumption are obtained by semi quantitative food frequency questionnaire (SQFFQ). Data about diabetic family history, ethnic, habit of smoking are obtained by interview. Data about physical activity is obtained by IPAQ questionnaire and data about obese an fasting blood glucose value are obtained by secondary data from GMC health center. Data were analyzed using chi square.

Results: There was significant association between diabetic family history, fried food's consumption frequency and the incidence of type 2 diabetes mellitus ($p < 0.05$). There was no significant association between ethnic, physical activity, habit of smoking, obese, eating pattern, fried food consumption (fried foods mass, fried foods energy and fried foods fat) ($p > 0.05$).

Conclusions: Risk factors related with type 2 DM evidence are diabetic family history and fried food's consumption frequency.

Keywords: Diabetes mellitus, Fried food consumption, Physical activity

INTRODUCTION

Degenerative disease is an epidemiologist transition from infectious diseases which dominated in some developing countries, including Indonesia. Type 2 diabetes mellitus (DM) is one of the disease with increasing prevalence

year by year. WHO predicted there are 300 million people with type 2 diabetes mellitus in 2025.

Health basic survey of Indonesia in 2013 showed that highest diabetes cases is in Yogyakarta (2.6%).¹ According to GadjahMada Medical Center (GMC), when diagnosed Universitas Gadjah Mada staffs, there are signs

and symptoms related to metabolic disorder, such as hypercholesterolemia, cardiovascular disease risk factor, and also type 2 DM.

The risk factor of type 2 DM related to the society's preference about fast food containing high fat and protein with low complex carbohydrate, especially fiber.² Moreover, obesity has greater risk up to 5.5 times to get type 2 DM than normal people.³ People with central obesity has tendency to get type 2 DM 3.9 times greater than normal people. Family history of type 2 DM also has relationship with type 2 DM as much as 5.5 times greater. Moreover, race or ethnic and physical activity also contribute to type 2 DM prevalence.^{4,6}

¹⁹ This study aims to observe the risk factors of type 2 DM. This finding would be an alternative intervention to prevent more patients of type 2 DM. Moreover, this finding can prevent the complication risk factors from type 2 DM.

²¹ **METHODS**

This is an observational cross sectional study. Research happened in GadjahMada Medical Center (GMC), Yogyakarta in March and April 2014. The study population is UGM staffs registered as member in GMC. The inclusion criteria are: UGM staffs regularly have medical checked up there, well communicated and has good memory, tend to be observed and interviewed, and also signed the inform consent. There are 179 persons as sample.

Data were collected and classified as: respondent identity, diagnosed has type 2 DM, race, smoking habit which collected by interview, physical activity collected with IPAQ, BMI, and also fasting blood glucose got from secondary data of GMC data, and food pattern, fried food consumption collected with *semi quantitative frequency questionnaire* (SQFFQ).

Sample categorized based on family history. It is divided into "yes group" who have minimal 1 family member with type 2 DM and "no group" if there is none. Race is divided into 2, Javanese and non-Javanese. Physical activity divided into 3 groups, high, moderate, and light based on IPAG category.⁷ Smoking habit divided into 2 groups, not smoking, and smoking more than 1 cigarette per day. Category of obesity is someone with BMI ≥ 25 kg/m².⁸ Type 2 DM diagnosed by fasting blood glucose as much as ≥ 126 mg/dl.⁹ Food pattern were categorized into 3 groups, high (>110% Indonesian recommended dietary intake (AKG)), well (90-110% AKG), and also low (<90% AKG). Fried food consumption were asked about consumption frequency, amount, energy and fat value of the food. Consumption frequency was categorized into 2 groups, frequent (≥ 4 x/week) and rare (<4x/week). Amount, energy, and fat value were divided into high (>mean/ median) and low (<mean/median).

Mean value used if data was normally distributed. Median value used if it is not. Data were analyzed with chi square test.

RESULTS

Sample's characteristic: Based on Table 1, more than half of the sample was 40-50 years old (50.8%), male (67%), had high education level (57%), and never been diagnosed type 2 DM (91.1%).

Table 1: Basic characteristic of sample in GMC.

Characteristic	Frequency n=179	Percentage (%)
Age group (years)		
40-50	91	50.8
≥ 50	88	49.2
Sex		
Male	120	67.0
Female	59	33.0
Education level		
High	102	57.0
Moderate	72	40.2
Low	5	2.8
Diagnosed type 2 DM		
Yes	16	8.9
No	163	91.1

High education level: finished university; Moderate education level: finished high school; Low education level: finished junior high school, elementary school.

Chi square table result was shown in Table 2, there was significant result according to age group and type 2 DM accident ($p > 0.05$), with RP = 0.197 (CI 95%: 0.054-0.716). Significant result also shown from education level with type 2 DM ($p < 0.05$) with RP=5.955 (CI 95%: 1.314-27.095). In addition, occupation as lecturer also has a relationship with type 2 DM accident ($p < 0.05$) with RP=3.17 (CI 95%: 1.117-9.016). On the other hand, there was no significant result between sex and type 2 DM accident ($p > 0.05$) with RP=1.475 (CI 95%: 0.471-4.958).

Risk factor of type 2 DM accident: The risk factors were shown in Table 3. Most of the subjects don't have type 2 DM in their family history (84.9%), ethnically Javanese (97.2%), never been smoked (85.5%), moderate physical activity (50.3%), non-obese (60.3%), low energy intake (70.9%), low fat intake (73.2), low intake protein (48%), low carbohydrate intake (71.5%), high fiber intake (63.7%), rarely eat fried food (55.9%), small amount of consumed fried food (50.3%), and also low fat proportion from consumed fried food (50.3%).

⁴ Table 4 showed that there is a significant relationship between heredity factors and type 2 DM accident ($p > 0.05$) with RP=0.132 (CI 95%: 0.044-0.393). In addition, there is also a significant relationship between frequency of consuming fried food with type 2 DM accident ($p < 0.05$) with RP=0.325 (CI 95%: 0.108-0.979).

In contrast, there is no significant relationship among ethnic, smoking habit, physical activity, obese, energy intake, fat intake, protein intake, carbohydrate intake, fiber intake, amount of the fried food, energy from fried food, and fat from fried food with type 2 DM accident (p>0.05).

Table 2: Relationship between respondent's characteristic and type 2 DM accident.

	Diabetes mellitus				Total		p	RP CI (95%)
	Yes n	%	No n	%	n	%		
Age group								
40-50	3	1.7	88	49.2	91	50.8	0.007*	0.197 (0.054-0.716)
≥50	13	7.3	75	41.9	88	49.2		
Sex							0.478	1.475 (0.471-4.958)
Male	12	6.7	108	60.3	120	67.0		
Female	4	2.2	55	30.7	59	33.0		
Education level							0.010*	5.966 (1.314-27.095)
High	14	7.8	88	49.2	102	57.0		
Moderate and low	2	1.1	75	41.9	77	43.0		
Lecturer							0.024*	3.173 (1.117-9.016)
Yes	9	5.0	47	26.3	56	31.3		
No	7	3.9	116	64.8	123	68.7		

Table 3: Risk factors of type 2 DM.

Variables		Frequency (n=179)	Percentage (%)
Family history	Yes	27	15.1
	No	152	84.9
Ethnic	Javanese	174	97.2
	Non-javanese	5	2.8
Smoking	Yes	26	14.5
	No	153	85.5
Physical activity	High	31	17.3
	Moderate	90	50.3
	Low	58	32.4
Obese	Yes	71	39.7
	No	108	60.3
Energy intake	High	19	10.6
	Adequate	33	18.4
	Low	127	70.9
Fat intake	High	24	13.4
	Adequate	24	13.4
	Low	131	73.2
Protein intake	High	52	29.1
	Adequate	41	22.9
	Low	86	48.0
Carbohydrate intake	High	17	9.5
	Adequate	34	19.0
	Low	128	71.5
Fiber intake	High	114	63.7
	Adequate	5	2.8
	Low	60	33.5
Frequency consume fried food	Frequent	79	44.1
	Rare	100	55.9
Amount of fried food	High	89	49.7
	Low	90	50.3
Fried food's amount of energy	High	88	49.2
	Low	91	50.8
Fried food's amount of fat	High	89	49.7
	Low	90	50.3

Table 4: The relationship of risk factors with type 2 DM accident.

	Diabetes mellitus				Total		p	RP CI (95%)
	Yes		No		n	%		
	n	%	n	%				
Family history								
Yes	8	4.5	19	10.6	27	15.1	0.001*	0.132 (0.044-0.393)
No	8	4.5	144	80.4	152	84.9		
Ethnic								
Javanese	16	8.9	158	88.3	174	97.2	0.623	0.908 (0.866-0.952)
Non-Javanese	0	0	5	4.6	5	2.8		
Smoking habit								
Yes	1	0.6	25	14.0	26	14.5	0.289	2.717 (0.343-21.504)
No	15	8.4	138	77.1	153	85.5		
Physical activity								
High	3	1.7	28	15.6	31	17.3	0.549	1.113 (0.297-4.164)
Moderate+low	13	7.3	135	75.4	148	82.7		
Obese								
Yes	9	5.0	62	98.3	71	39.7	0.155	0.477 (0.169-1.347)
No	7	3.9	101	56.4	108	60.3		
Energy intake								
High	1	0.6	18	10.1	19	10.6	0.472	0.537 (0.067-4.310)
Adequate+low	15	8.9	145	81.0	160	89.4		
Fat intake								
High	1	0.6	23	12.8	24	13.4	0.335	0.406 (0.051-3.221)
Adequate+low	15	8.9	140	78.2	155	86.6		
protein intake								
High	5	2.8	47	26.3	52	29.1	0.520	1.122 (0.370-3.405)
Adequate+low	11	6.1	116	64.8	127	70.9		
carbohydrate intake								
High	2	1.1	15	8.4	17	9.5	0.464	1.410 (0.292-6.802)
Adequate+low	14	7.8	148	82.7	162	90.5		
Asupan Serat								
Tinggi	12	6.7	102	57.0	114	63.7	0.242	1.794 (0.554-5.811)
Baik+Rendah	4	2.2	61	34.1	65	36.3		
Frequency consume fried food								
Frequent	11	6.1	68	38.0	79	44.1	0.038*	0.325 (0.108-0.979)
Rare	5	2.8	95	53.1	100	55.9		
Amount of fried food								
High	7	3.9	82	45.8	89	49.7	0.617	1.302 (0.463-3.662)
Low	9	5.0	81	45.3	90	50.3		
Energy from fried food								
High	8	4.5	80	44.7	88	49.2	0.944	0.964 (0.345-2.692)
Low	8	4.5	83	46.4	91	50.8		
Fat from fried food								
High	6	3.4	83	46.4	89	49.7	0.306	1.729 (0.600-4.979)
Low	10	5.6	80	44.7	90	50.3		

DISCUSSION

The relationship between sample's characteristic with diabetes mellitus

¹⁷ There is a significant result between age group with type 2 DM accident among the subject ($p < 0.05$) and the value ¹³ RP is 0.197 (CI 95%: 0.054-0.716). Approximately 54.33% of type 2 diabetes has been observed in the age group >40-50 years.¹⁰ It has been known that people

⁴ above 45 years old are tend to have higher risk of type 2 DM and glucose intolerance. It is caused by degenerative factor that reduce physical ability to metabolite glucose. Though that's not only caused by age but also by their abilities to survive.¹¹

There is also a significant relationship between education level and type 2 DM accident ($p < 0.05$) where the RP is 5.955 (CI 95%: 1.314-27.095). This result is similar to Yunita's in GMC. She found that education level is significantly related to metabolic syndrome.¹² Education

level is related to people's knowledge leads to their food preference and life style, which affect to degenerative syndrome, such as type 2 DM.

The risk factors of diabetes mellitus

Heredity

Statistic test showed a significant result ($p < 0.05$) with $RP = 0.132$ (CI 95%: 0.044-0.393) between heredity and type 2 DM among the subject. Family history has a strong possibility to affect the accident of type 2 DM, the OR is 3.75.¹³ From this we can conclude that people came from diabetic family are willing to get this disease 3.75 times greater than others. Research conducted in Sweden found that mutation of hepatocyte nuclear factor (HFN)-1 α and (HFN)-4 α provoke type 2 DM.¹⁴

Ethnic

Ethnic is not significantly related to type 2 DM ($p > 0.05$) with $RP = 0.908$ (CI 95%: 0.866-0.952). It may be caused by an equal proportion between Javanese respondents and non-Javanese.

Food pattern is composed by human needs and culture called biocultural interfece.¹⁵ Moreover, another factors affect food pattern are ecology, social environment, technology, and culture it self.¹⁶ Cultural research showed that most of *Minang* people are willing to have spicy and curry food. On the other hand, Javanese people love sweet food. It is not absolute for all the people, but there is similar mind in society about their own food preference.¹⁷

Smoking habit

The relationship between smoking habit and type 2 DM is not significant ($p > 0.05$), $RP = 2.717$ (CI 95%: 0.343-21.504). It can be concluded that smoking habit is not the risk factor. Smoking habit is strongly related to type 2 DM ($p < 0.05$), OR 2.6.¹⁸ Smoking habit is not the predictor that caused death in people with type 2 DM who also have CVD.¹⁹

23 *Physical activity*

The relationship between physical activity and type 2 DM is not significant ($p > 0.05$) with the value of $RP = 1.113$ (CI 95%: 0.297-4.164). Another study conducted in Riau found that physical activity is not significantly related to glucose blood value ($p = 0.385$).²⁰ On the other hand, another **16**earch found that physical activity is related to type 2 DM ($p = 0.024$ dan OR 2.37).²¹ Physical activity can increase blood circulation, reduce body weight, and repair insulin sensitivity, which lead to better glucose blood level.²¹ The different result may be caused by the major characteristic of the subject have moderate level of physical activity. Moreover, type 2 DM is caused by a lot

of factors, whereas food pattern and physical activity are not the only one.

Obese

Obese is not a significant risk factor of type 2 DM ($p > 0.05$), $RP = 0.477$ (CI 95%: 0.169-1.347). Another study found that obesity is related to type 2 DM ($p < 0.05$), $OR = 1.66$ (CI 95%: 1.35-2.03).²² Another study found that 65.5% are centrally obese.²³ Another study also found that obesity is not significantly related to type 2 DM ($p = 0.435$).²⁴

Overweight people has higher leptin level in body. Leptin is a hormone related to gene that caused obesity. In hipotalamus, leptin regulates lipid level and converts fat into energy. Leptin works in peripheral nervous system and central system. Leptin can inhibit insulin receptorsubstrate-1(IRS). It reduces intake glucose in cell, so the glucose blood level is increased.²⁵ Fat people is not always have type 2 DM, this study found that obesity is not the key factor of type 2 DM. This disease caused by sedentary lifestyle.²⁶

Food pattern

Food pattern with high energy is not significantly related to type 2 DM ($p > 0.05$), $RP = 0.537$ (CI 95%: 0.067-4.310). On the other hand, Lasmia found that energy intake related to type 2 DM ($p < 0.05$, $OR = 4.8$ CI 95%: 1.8-12.4).²⁷ Excess of high energy intake caused insulin resistant. Glucose and free fatty acid in blood are excessive and adipose tissue in body also increase and caused obesity. Central obesity is strongly known related to insulin resistant.²⁸

Fat intake also not significantly related to type 2 DM ($p < 0.05$, $RP = 0.406$, CI 95%: 0.05-221). Juleka in her study said that fat intake affects blood glucose level in people with type 2 DM ($p < 0.05$, $OR = 5.20$, CI 95% 1.08-24.89).²⁹ Excessive fat intake makes more deposit energy storage. Moreover, fat inhibit gene produced mitochondria. It leads to broken metabolism of cell, including in insulin response. In conclusion, people consumed high fat food are tend to have type 2 DM.³⁰

Amount of protein consumption also not significantly related to type 2 DM ($p > 0.05$, $RP = 1.122$, CI 95%: 0.370-3.405). Another study also has similar result that protein is not the risk factor of type 2 DM ($p > 0.05$, CI 95%: 1.0-6.2).²⁷ Carbohydrate consumption also not significantly related **8** type 2 DM ($p > 0.05$, $RP = 1.41$, CI 95%: 0.292-6.802). On the other hand, a study found that type 2 DM is related to carbohydrate intake ($p < 0.05$, $OR = 4.67$, CI 95%: 1.35-16.19).³⁰ Reducing carbohydrate consumption can increase insulin sensitivity in healthy people, which leads to lower glucose blood in diabetic people.²⁸

Fiber consumption has no significant relationship with type 2 DM ($p > 0.05$, $RP = 1.794$, CI 95%: 0.554-5.811). In

contrast, a study found that fiber intake is strongly related to type 2 DM.²⁸ Higher fiber intake leads to slower digestive speed from gastric to intestine. It inhibit fast increasing in post-prandial blood glucose.²⁹

Some results are different with the previous study and theories. It may be caused by the characteristic of the subject. Most of them have low energy, carbohydrate, protein, and fiber intake. Moreover, there isn't any robust different between healthy people's intake and diabetic people's.

Konsumsi gorengan

Fried food consumption is significantly related to type 2 DM ($p < 0.05$, $RP = 0.325$, CI 95%: 0.108-0.979). The amount of the fried food ($p > 0.05$, $RP = 1.302$, CI 95%: 0.463-3.662) and the amount of energy in fried food are not relevant with type 2 DM accident ($p > 0.05$, $RP = 1.729$, CI 95%: 0.600-4.979). Fried food is a high fat food. People with type 2 DM are tend to get heart and vascular disease. This is the strong reason why fat and cholesterol should be consume in a small portion.³¹

CONCLUSION

Heredity and the frequency of fried food consumption are the two risk factors of type 2 DM. We wish that people nowadays have better food preference and avoid trans and saturated fat. For further study, the relationship among low sugar food, low G5 food, soluble fiber and reducing in glucose blood level should be conducted.

Funding: No funding sources

Conflict of interest: None declared

Ethical approval: The study was approved by the Institutional Ethics Committee

REFERENCES

1. Kemenkes. Riset Kesehatan Dasar: Riskesdas 2013. Badan Penelitian dan Pengembangan Kesehatan Kementerian Kesehatan RI. 2013.
2. Sugianti, L. Hubungan antara Konsumsi Kafein dengan Sindrom Metabolik pada Peserta GMC Health Center. Skripsi. Universitas Gadjah Mada. 2010.
3. Husain SA, Pangastuti R. (2010). Pengendalian Status Gizi, Kadar Glukosa Darah dan Tekanan Darah Melalui Terapi Gizi Medis pada Pasien Diabetes Mellitus (DM) Tipe 2 Rawat Jalan di RSU Mataram NTB. Indonesian J Clinical Nutrition. 2010;7(2):28-57.
4. Wiardini, Ni Komang. (2007). Pola Makan dan Obesitas sebagai Faktor risiko DM tipe II di RS Sanglah Denpasar. Jurnal Gizi Klinik Indonesia. 2007;4(1):1-10.
5. Fauci AS, Kasper DL, Longo DL, Braunwald E, Hauser SL, Jameson JL, Loscalzo J. Harrison's

- Principles of Internal Medicine. 7th edition. New York : Mc Graw Hill. 2008.
6. Norris SL, Kansagara, D, Bougatsos C, Nygren P, Fu R. Screening for Type 2 Diabetes Mellitus: Update of 2003 Systematic Evidence Review for the U.S. Preventive Services Task Force. Agency for Health care Research and Quality. 2008;1-3.
7. IPAQ. Guidelines for Data Processing and analysis of the International Physical Activity Questionnaire (IPAQ) Short Form. 2005. Available from: www.ipaq.ki.se[diakses 14 Desember 2013].
8. Inoue S, Zimmet P, Caterson I, Chunming C, Ikeda, Y, Khalid AK. The Asia Pasific Perspectiv: Redefining Obesity and its Treatmen, Health Communication Australia Pty Limited. 2000.
9. Perkeni, Perkumpulan Endokrinologi Indonesia. Konsensus Pengendalian dan Pencegahan Diabetes Mellitus Tipe2 di Indonesia 2011. Jakarta.
10. Nandimath VA, Swamy CS, Nandimath SA, Jatti, G, Jadhav S. Evaluation of certain risk factors of type 2 diabetes mellitus: a case-control study. Int J Med Sci Public Health. 2016;5(7): 1334-9.
11. Wicaksono, Putro R. Faktor-Faktor yang Berhubungan dengan Kejadian Diabetes Mellitus Tipe 2. Skripsi. Universitas Diponegoro. 2012.
12. Riska Y. Hubungan Pola Aktivitas Fisik dengan Sindroma Metabolik pada Peserta GMC Health Center. Skripsi. Universitas Gadjah Mada. 2012.
13. Zahtamal, Chandra F, Suyanto, Restuastuti T. Faktor-Faktor Risiko Pasien Diabetes Mellitus. Berita Kedokteran Masyarakat. 2007;23(3):142-7.
14. Holmkvist J, Almgren P, Lyssenko V, Lindgren, CM, Eriksson KF, Isomaa B, Tuomi T, Nilsson P, Groop L. Common Variants in Maturity-Onset Diabeteses of the Young Genes and Future Risk of type 2 Diabetes. In: Diabetes. 2006;57.
15. Sanjur D. Social and Cultural Perapektifes in Nutrition. Washington DC: Prentice Hall, Inc. New York, USA. 1982.
16. Jerome, NW, Peltó GH, Kandel RF. An Ecological Approach to Nutritional Anthropology. New York : Redgrave. 1980.
17. Ariani NI. (2013). Strategi Adaptasi Orang Minang terhadap Bahasa, Makanan dan Norma Masyarakat Jawa. Jurnal Komunitas Unnes. 2013;(1):26-37.
18. Mujio. Faktor-Faktor Risiko Kejadian Penyakit Diabetes Mellitus Tipe 2 pada Orang Dewasa di Kabupaten Boyolali Propinsi Jawa Tengah. Tesis. Universitas Gadjah Mada. 2006.
19. Muggeo M, Zoppini G, Brun E, Bonora E, Verlató G. Mortality and Its Predictors In Type 2 Diabetes. In: Sinclair, A.J, Finucane, P, Diabetes in Old Age, New York, Jonh Wiley & Sons, LTD. 2001.
20. Chandra F, Masdar H, Rosdiana D. Identifikasi Pola Aktivitas Fisik dan Status Gizi Pegawai Negeri Sipil Pemerintah Daerah Provinsi Riau dan hubungan dengan kadar gula darah. Karya Ilmiah. Fakultas Kedokteran Universitas Riau. 2013.
21. Manik HR. Faktor Risiko yang Bisa Dimodifikasi Terhadap Diabetes Mellitus Tipe 2 di RSU

- Hadrianus Sinaga Pangururan Kabupaten Samosir. Tesis. Universitas Sumatra Utara. 2012.
22. Soetiarto F, Roselinda, Suhardi. Hubungan Obesitas dengan Diabetes Mellitus berdasarkan IMT dan Lingkar Pinggang Data Riskesdas 2007. Buletin Penelitian Kesehatan. 2010;38(1):36-42.
 23. Anjana P, Bant DD. A community based cross-sectional study to identify individuals at high risk for diabetes in urban slums of Hubli. *Int J Community Med Public Health*. 2016;3(1): 212-7
 24. Hikmat R. Hubungan Obesitas dengan Diabetes Mellitus di Poliklinik Penyakit Dalam RSUD Cibabat Cimahi Tahun 2010. *Jurnal Kesehatan Kartika*. hal: 2010;7-13.
 25. Peter DJ. Diet Sehat Diabetes sesuai Golongan Darah. Yogyakarta: Delapratasa. 2008.
 26. Misnadiarly. Obesitas Sebagai Faktor Resiko Beberapa Penyakit Jantung. Jakarta: Pustaka Obor Populer. 2007.
 27. Lasmiasi. Pola Makan sebagai Faktor Risiko Terhadap Kejadian Diabetes Mellitus tipe 2 pada RSUD di Jayapura. Skripsi. Universitas Gadjah Mada. 2009.
 28. Fitri RI, Yekti W. Asupan Energi, Karbohidrat, Serat, Beban Glikemik, Latihan Jasmani dan Kadar Gula Puasa pada Pasien Diabetes Mellitus Tipe 2. *Jurnal Media Medika Indonesiana*. 2012;46(2):121-31.
 29. Juleka. Hubungan Pola Makan dengan Pengendalian Kadar Glukosa Darah Pengidap Diabetes Mellitus Tipe 2 Rawat Jalan di RSUD Gunung Jadi Cirebon. Tesis. Universitas Gadjah Mada. 2005.
 30. Kartini. Hubungan Pola Konsumsi Makanan dengan Kejadian Diabetes Mellitus Tipe 2 Pada Penduduk Wanita di Kota Banjarmasin. Tesis. Universitas Gadjah Mada. 2012.
 31. Rahmawati. Pola Makan dan Aktifitas Disik dengan Kadar Glukosa Darah Penderita Diabetes Mellitus tipe 2 Rawat Jalan di RSUD Dr. Wahidin Sudirohusodo Makasar. Skripsi. Universitas Hasanuddin. 2011.

Cite this article as: Sudargo T, Pertiwi S, Alexander RA, Siswati T, Ernawati Y. The relationship between fried food consumption and physical activity with diabetes mellitus in Yogyakarta, Indonesia. *Int J Community Med Public Health* 2017;4:38-44.

The relationship between fried food consumption and physical activity with diabetes mellitus in Yogyakarta, Indonesia

ORIGINALITY REPORT

13%

SIMILARITY INDEX

12%

INTERNET SOURCES

9%

PUBLICATIONS

7%

STUDENT PAPERS

PRIMARY SOURCES

1	eprints.arums.ac.ir Internet Source	3%
2	Submitted to Aspen University Student Paper	2%
3	pdfs.semanticscholar.org Internet Source	2%
4	scifes.fkm.ui.ac.id Internet Source	1%
5	mail.ijsurgery.com Internet Source	1%
6	R. M. Stoney. "Insulin resistance as a major determinant of increased coronary heart disease risk in postmenopausal women with Type 2 diabetes mellitus", Diabetic Medicine, 6/2001 Publication	1%
7	repository.usu.ac.id Internet Source	1%

8	<p>Mahalul Azam, Fetty Nur Hidayati, Arulita Ika Fibriana, Udin Bahrudin, Syed Mohamed Aljunid. "Prevalence of Isolated Systolic Hypertension among People with Diabetes in Indonesia", Cold Spring Harbor Laboratory, 2020</p>	1 %
Publication		
9	<p>ejournal.undip.ac.id</p>	<1 %
Internet Source		
10	<p>mafiadoc.com</p>	<1 %
Internet Source		
11	<p>www.ajol.info</p>	<1 %
Internet Source		
12	<p>Sung Keun Park, Mi Hae Seo, Ho Cheol Shin, Jae-Hong Ryoo. "Clinical availability of nonalcoholic fatty liver disease as an early predictor of type 2 diabetes mellitus in korean men: 5-year prospective cohort study", Hepatology, 2013</p>	<1 %
Publication		
13	<p>www.ejmanager.com</p>	<1 %
Internet Source		
14	<p>worldwidescience.org</p>	<1 %
Internet Source		
15	<p>www.worldcocoafoundation.org</p>	<1 %
Internet Source		

16	123dok.com Internet Source	<1 %
17	jurnal.syedzasaintika.ac.id Internet Source	<1 %
18	qdoc.tips Internet Source	<1 %
19	ujpr.org Internet Source	<1 %
20	www.frontiersin.org Internet Source	<1 %
21	"RHEUMATOID ARTHRITIS ? TREATMENT", APLAR Journal of Rheumatology, 8/2006 Publication	<1 %
22	Mia Audina, Tonny Cortis Maigoda, Tetes Wahyu W. "Status Gizi, Aktivitas Fisik dan Asupan Serat Berhubungan dengan Kadar Gula Darah Puasa Penderita DM Tipe 2", Jurnal Ilmu dan Teknologi Kesehatan, 2018 Publication	<1 %
23	dx.doi.org Internet Source	<1 %

Exclude quotes Off
Exclude bibliography On

Exclude matches < 7 words