

DAFTAR PUSTAKA

- A., A. and T., B. (2016) ‘Incidence of lower limb motor weakness in patients receiving postoperative epidural analgesia and factors associated with it: An observational study’, *Saudi Journal of Anaesthesia*, pp. 149–153. Available at: <http://www.embase.com/search/results?subaction=viewrecord&from=export&id=L608962243%0A> <http://dx.doi.org/10.4103/1658-354X.168806>.
- Amalia, F. Y. (2020) ‘Hubungan Dukungan Keluarga Dengan Pelaksanaan Mobilisasi Dini Pada Pasien Post Operasi Di Ruang Bedah Rsud Dr . H . Bob Bazar , SKM’, *Jurnal Ilmu Kesehatan Indonesia (JIKSI)*, 1(1), pp. 1–8.
- Amornyotin, S. (2016) ‘Anesthetic Consideration for Geriatric Patients’, *Intech*, i(tourism), p. 13.
- Asgari, Z. et al. (2017) ‘Spinal Anesthesia and Spinal Anesthesia with Subdiaphragmatic Lidocaine in Shoulder Pain Reduction for Gynecological Laparoscopic Surgery: A Randomized Clinical Trial’, *Pain Research and Management*, 2017. doi: 10.1155/2017/1721460.
- Ayuningtyas1, A. F., Susilo2, C. B. and Ana Ratnawati3 (2020) ‘FACTORS RELATED TO ACHIEVING BROMAGE SCORE IN PATIENTS’, pp. 2–3.
- Bothe, J. (2017) ‘Postoperative Mobilisation Guideline’.
- Breebaart, M. B. (2019) ‘Local Anaesthetics for Spinal Anaesthesia in Day-Case Surgery’, *Intech*, i(tourism), p. 13.
- Cakir, H. et al. (2013) ‘Adherence to enhanced recovery after surgery and length of stay after colonic resection’, *Colorectal Disease*, 15(8), pp. 1019–1025. doi: 10.1111/codi.12200.
- Capdevila, X. et al. (2020) ‘Factors Determining the Choice of Spinal Versus General Anesthesia in Patients Undergoing Ambulatory Surgery: Results of a Multicenter Observational Study’, *Advances in Therapy*, 37(1), pp. 527–540. doi: 10.1007/s12325-019-01171-6.
- Chen, C. P. C. et al. (2021) ‘The lateral decubitus body position might improve the safety of ultrasound-guided supraclavicular brachial plexus nerve block’, *Journal of Pain Research*, 14, pp. 75–82. doi: 10.2147/JPR.S276095.
- Cheng, H. et al. (2018) ‘Prolonged operative duration is associated with complications: a systematic review and meta-analysis’, *Journal of*

- Surgical Research*, 229, pp. 134–144. doi: 10.1016/j.jss.2018.03.022.
- Elis Roslianti, Yanti Srinayanti, Jajuk Kusumawaty, E. Setyani (2020) ‘the Description of the Treatment of Early Mobilization on the Mother Post Sectio Caesarea in Lotus II Blud Hospitals Banjar Year 2018’, 7(April), pp. 49–62.
- Fabian Grass, Basile Pache, David Martin, Valérie Addor, Dieter Hahnloser, Nicolas Demartines, M. H. (2018) ‘Feasibility of early postoperative mobilisation after colorectal surgery: A retrospective cohort study’. *Feasibility of early postoperative mobilisation after colorectal surgery: A retrospective cohort study*.
- Faculty of pain medicine, T. R. C. of A. (2020) ‘Best Practice in the Management of Epidural Analgesia in the Hospital Setting’, (August), pp. 1–18.
- Fahruruddin, Amri, I. and Wahyudi (2017) ‘Perbandingan Efek Antara Dexmedetomidin Dosis 0.25 mcg/kgBB Dan 0.5 mcg/kgBB Intravena Terhadap Durasi Blok Anestesi Spinal Pada Bedah Ekstremitas Bawah’, *Jurnal Kesehatan Tadulako*, 3(2), pp. 9–20.
- Fleisher, L. A. (2018) ‘Regional Anesthesia: What We Need to Know in the Era of Enhanced Recovery After Surgery Protocols and the Opioid Epidemic’, *Anesthesiology Clinics*, 36(3), pp. xi–xii. doi: 10.1016/j.anclin.2018.06.002.
- Frayoga, F. and Nurhayati, N. (2018) ‘Pengaruh Mobilisasi Dini terhadap Pemulihan Kandung Kemih Pasca Pembedahan dengan Anestesi Spinal’, *Jurnal Ilmiah Keperawatan Sai Betik*, 13(2), p. 226. doi: 10.26630/jkep.v13i2.936.
- Garmon, D. J. D. A. G. P. B. E. H. (2021) ‘American Society of Anesthesiologists Classification - StatPearls - NCBI Bookshelf’. Available at: <https://www.ncbi.nlm.nih.gov/books/NBK441940/>.
- Hadbi Mohamed*, Benalakma Dalila, Ramzi Berbiche, Fatma Bouderbala, F. N. (2021) ‘Journal of Anesthesia & Clinical Abnormal Prolonged Duration of Spinal Anesthesia in Patient with Multiple Sclerosis’, pp. 1–4.
- Hamzeh Hosseinzadeh, Dawood Aghamohammadi, Marzieh Marahem, Negin Magsumi, P. H. (2019) ‘Corellation of Body Mass Index and Spread of Spinal Block in Herniorraphy Patients under Spinal Anesthesia’, 4(October), pp. 3–6.
- Hunie, M. et al. (2021) ‘The current practice of spinal anesthesia in anesthetists at a comprehensive specialized hospital: A single center observational study’, *Local and Regional Anesthesia*, 14, pp. 51–56. doi:

10.2147/LRA.S300054.

- Jeffrey M. Carness and Mark J. Lenart (2013) ‘Current Local Anesthetic Applications in Regional Anesthesia’, *InTech*, 32(July), pp. 137–144. Available at: <http://www.intechopen.com/books/trends-in-telecommunications-technologies/gps-total-electron-content-tec-prediction-at-ionosphere-layer-over-the-equatorial-region%0AInTec%0Ahttp://www.asociatiamhc.ro/wp-content/uploads/2013/11/Guide-to-Hydropower.pdf>.
- Kaye, A. D., Urman, R. D. and Vadivelu, N. (2012) *Essentials of regional anesthesia, Essentials of Regional Anesthesia*. doi: 10.1007/978-1-4614-1013-3.
- Keller, D. S. and Chand, M. (2017) ‘Increasing patient engagement to move enhanced recovery forward’, *Annals of Laparoscopic and Endoscopic Surgery*, 2, pp. 155–155. doi: 10.21037/ales.2017.10.03.
- Leger, R., Livesberger, J. and Sinha, A. (2020) ‘Enhanced recovery after surgery (ERAS) in clinical practice’, *Anaesthesia, Pain and Intensive Care*, 24(3), pp. 335–345. doi: 10.35975/apic.v24i3.1287.
- Ljungqvist, O., Scott, M. and Fearon, K. C. (2017) ‘Enhanced recovery after surgery a review’, *JAMA Surgery*, pp. 292–298. doi: 10.1001/jamasurg.2016.4952.
- Mahan, M. C. et al. (2019) ‘Time of return of neurologic function after spinal anesthesia for total knee arthroplasty: mepivacaine vs bupivacaine in a randomized controlled trial’, *Arthroplasty Today*, 5(2), pp. 226–233. doi: 10.1016/j.artd.2019.03.003.
- Matthews, C. and Hospital, D. (2010) ‘Enhanced Recovery After Surgery (Eras) Anaesthesia Tutorial of the Week 204 !’, *Atotw*, (November), pp. 1–9.
- Matthews, H. et al. (2010) ‘Early Mobilization in Trauma ICU Decreases Length of Stay and Complications.’, *Journal of Acute Care Physical Therapy*, p. 72. doi: 10.1097/01592394-201001020-00010.
- Mikhail’s, M. & (2013) *Clinical Anesthesiology, Saudi Journal of Anaesthesia*. doi: 10.4103/1658-354X.109819.
- Moningi, S. et al. (2019) ‘Enhanced recovery after surgery: An anesthesiologist’s perspective’, *Journal of Anaesthesiology Clinical Pharmacology*, pp. 5–13. doi: 10.4103/joacp.JOACP_238_16.
- Olawin, A. J. M. (2021) ‘Spinal Anesthesia - StatPearls - NCBI Bookshelf’, *StatPearl Publishing*. Available at: <https://www.ncbi.nlm.nih.gov/books/NBK537299/>.

- Pasternak, S., Schwab, N. and Thun, V. (2018) 'Implementation of an enhanced recovery after surgery program', *Nursing Critical Care*, 13(4), pp. 40–45. doi: 10.1097/01.CCN.0000532368.48473.17.
- Pederson, J. L. et al. (2020) 'The impact of delayed mobilization on postdischarge outcomes after emergency abdominal surgery: A prospective cohort study in older patients', *PLoS ONE*, 15(11 November), pp. 1–13. doi: 10.1371/journal.pone.0241554.
- Rustianawati, Y. et al. (2013) 'Efektivitas Ambulasi Dini terhadap Penurunan Intensitas Nyeri pada Pasien Post Operasi Laparotomi di RSUD Kudus', *Jurnal Ilmu Keperawatan dan Kebidanan (JIKK)*, pp. 1–8. Available at: <https://ejr.stikesmuhkudus.ac.id/index.php/jikk/article/download/203/134>.
- Shergill, S. and Galway, U. (2018) 'Atypical prolonged spinal anaesthesia', *Indian Journal of Anaesthesia*, pp. 1004–1005. doi: 10.4103/ija.IJA_527_18.
- Sholihah 'Ush, A. (2017) 'Pengaruh Mobilisasi Dini Terhadap Peningkatan Kekuatan Otot Pada Pasien Pasca Stroke Iskemik Di Rsud Dr. Harjono Ponorogo', *Вестник Росздравнадзора*, 4, pp. 9–15.
- Da Silva Filho, A. L. et al. (2018) 'Enhanced Recovery After Surgery (ERAS): New concepts in the perioperative management of gynecologic surgery', *Revista Brasileira de Ginecologia e Obstetricia*, 40(8), pp. 433–436. doi: 10.1055/s-0038-1668581.
- Simin, A. et al. (2018) 'Effect of position during induction of spinal anaesthesia for caesarean section on maternal haemodynamic: Randomised clinical trial', *Journal of Clinical and Diagnostic Research*, 12(2), pp. UC05–UC08. doi: 10.7860/JCDR/2018/27753.11184.
- Soffin, E. M. and Yadeau, J. T. (2016) 'Enhanced recovery after surgery for primary hip and knee arthroplasty: A review of the evidence', *British Journal of Anaesthesia*, 117, pp. iii62–iii72. doi: 10.1093/bja/aew362.
- Sudha, H. (2016) 'Subarachnoid Block in Prone Position for Giant Gluteal Neurofibroma', *Journal of Anesthesia & Critical Care: Open Access*, 4(2), pp. 2–4. doi: 10.15406/jaccaa.2016.04.00131.
- Sudiharjani, N. (2012) 'Mobilisasi Dini Dan Penyembuhan Luka Operasi Pada Ibu Post Sectio Caesarea (Sc) Di Ruang Dahlia Rumah Sakit Umum Daerah Kota', *Prosiding Seminar Nasional*. Available at: <https://bem.unimus.ac.id/ojsunimus/index.php/psn12012010/article/view/1281>.

- Taeteti, A. M., Johan, A. and Anggorowati (2018) ‘Early Mobilisation Terhadap Pemulihan Pascaoperasi Laparotomi Di Rumah Sakit.’
- Tang, Z. *et al.* (2019) ‘Observation of single spinal anesthesia by 25G needle puncture through a lateral crypt for hip surgery in elderly patients’, *Medicine*, 98(27), p. e16334. doi: 10.1097/MD.00000000000016334.
- Twomey, R. *et al.* (2021) ‘Impact of early mobilization on recovery after major head and neck surgery with free flap reconstruction’, *Cancers*, 13(12), pp. 4–15. doi: 10.3390/cancers13122852.
- Upadhyay, R. K., Shenoy, L. and Venkateswaran, R. (2018) ‘Effect of intravenous dexmedetomidine administered as bolus or as bolus-plus-infusion on subarachnoid anesthesia with hyperbaric bupivacaine’, *Journal of Anaesthesiology Clinical Pharmacology*, 34(3), pp. 46–50. doi: 10.4103/joacp.JOACP.
- Zhang, L. *et al.* (2019) ‘Early mobilization of critically ill patients in the intensive care unit: A systematic review and meta-analysis’, *PLoS ONE*, 14(10), pp. 1–16. doi: 10.1371/journal.pone.0223185.