

DAFTAR PUSTAKA

- Asil, E., Cakiroglu, M. S., Pinar, S. F., Ucar, A., Ozcelik, A. O., Yilmaz, M. V., & Akan, L. S. (2014). Factors That Affect Body Mass Index of Adults. *Pakistan Journal of Nutrition*, 13: 255-26.
- Azmi, D. A., Wiyono, J., & Dtn, I. (2019). Relationship of Body Mass Index (BMI) and Type of Operation With Time of Conscious Recover in Postoperative Patients With General Anesthesia at Recovery Room of Bangil Hospital. *Jurnal Keperawatan Terapan (e-Journal)*, 05(02), 2442–6873.
- Bishop, D. G. (2014). Predicting Spinal Hypotension During Caesarean Section. *Southern African Journal of Anaesthesia and Analgesia*, 170–173. <https://doi.org/10.1080/22201181.2015.959336>
- Burhanuddin, Y. E., Syahrianti, & Afrianty, I. (2021). Perbedaan Kadar Interleukin 6 Serum dan Kadar HsCrp Pada Ibu Hamil Preeklampsia. *Jurnal Kesehatan*, Vol. 04.
- Butterworth, J. F., C, M. D., & Wasnick, J. D. (2018). *Morgan & Mikhail's clinical anaesthesiology*. New York : McGraw-Hill Education.
- Drasner, K., & Larson, M. (2011). *Basics of Anesthesia : Spinal and Epidural Anesthesia*. Elsevier.
- Fakherpour, A., Ghaem, H., & Zaree, Z. F. S. (2018). Maternal and anaesthesia-related risk factors and incidence of spinal anaesthesia-induced hypotension in elective caesarean section: A multinomial logistic regression. *Indian J Anaesth*, 62(1):36-46.
- Goel, K., Luthra, N., Goyal, N., Grewal, A., & Taneja, A. (2021). Comparison of norepinephrine and phenylephrine infusions for maintenance of haemodynamics following subarachnoid block in lower segment caesarean section. *Indian Journal of Anaesthesia*, 65(8), doi: 10.4103/ija.IJA_185_21.
- Hernández, M. G. L., Flórez, H. J. M., Robles, S. Á., & Arteaga, J. de L. A. (2018). Risk factors for hypotension in regional spinal anesthesia for cesarean section. Role of the Waist-to-Hip Ratio and Body Mass Index. *Colombian Journal of Anesthesiology*, 46(1), 42–48. <https://doi.org/https://doi.org/10.1097/cj9.0000000000000008>
- Huang, Y.-Y., & Chang, K.-Y. (2021). Sensory Block Level Prediction of Spinal Anaesthesia with 0.5% Hyperbaric Bupivacaine: a Retrospective Study. *Scientific Reports*, 9105.
- Latief, S. A., Suryadi, K. A., & Dachlan, M. R. (2015). *Petunjuk Praktis Anestesiologi*. Universitas Indonesia.

- Marashi, S. M., Soltani-Omid, S., Soltani Mohammadi, S., Aghajani, Y., & Movafegh, A. (2014). Comparing two different doses of intravenous ondansetron with placebo on attenuation of spinal-induced hypotension and shivering. *Anesthesiology and Pain Medicine Journal*, 4(2). <https://doi.org/10.5812/aapm.12055>
- Masturoh, I., & Anggita, N. (2018). *Metodologi Penelitian Kesehatan*. Pusat Pendidikan Sumber Daya Manusia dan Kesehatan.
- Mochtar, R. (2012). *Sinopsis Obstetri: Obstetri Fisiologi, Obstetri Patologi*. (Edisi ketiga), Jakarta : EGC.
- Nikooseresht, M., Rabiei, M. A. S., Hajian, P., Dastaran, R., & Alipour, N. (2016). Comparing the Hemodynamic Effects of Spinal Anesthesia in Preeclamptic and Healthy Parturients During Cesarean Section. *Anesthesiology and Pain Medicine*, 6(3):e11519. <https://doi.org/https://dx.doi.org/10.5812/aapm.11519>
- Notoatmodjo, S. (2012). *Metodologi Penelitian Kesehatan*. Jakarta: Rine Cipta.
- Nursalam. (2017). *Metodologi Penelitian Ilmu Keperawatan: Pendekatan Praktis*. Jakarta: Salemba Medika.
- Priyono. (2016). *Metode Penelitian Kuantitatif*. Sidoarjo: Zifatama Publishing.
- Pu, X., & Sun, J. (2019). General anesthesia vs spinal anesthesia for patients undergoing total-hip arthroplasty A meta-analysis. *Medicine*, 98:16, 4.
- Puspitasari, A. I. (2019). *Faktor-Faktor yang Mempengaruhi Kejadian Hipotensi pada Pasien dengan Spinal Anestesi di RSUP Dr. Soeradji Tirtonegoro Klaten*. <http://eprints.poltekkesjogja.ac.id/3571/1>.
- Puspitasari, N. (2018). Faktor Kejadian Obesitas Sentral pada Usia Dewasa. *Higeia Journal Of Public Health Research and Development*.
- Ranasinghe, C., Gamage, P., Katulanda, P., Andraweera, N., Thilakarathne, S., & Tharanga, P. (2013). *Relationship between Body mass index (BMI) and body fat percentage, estimated by bioelectrical impedance, in a group of Sri Lankan adults: A cross sectional study*. 13(1), BMC Public Health. <https://doi.org/10.1186/1471-2458-13-797>
- Rustini, R., Fuadi, I., & Surahman, E. (2016). *Insidensi dan Faktor Risiko Hipotensi pada Pasien yang Menjalani Seksio Sesarea dengan Anestesi Spinal di Rumah Sakit Dr. Hasan Sadikin Bandung*. 4(1), 42–49.
- Sastroasmoro, S., & Ismael, S. (2014). *Dasar dasar Metodologi Penelitian Klinis* (5). Jakarta : Sagung Seto.

- Šklebar, I., Bujas, T., & Habek, D. (2019). Spinal Anaesthesia-Induced Hypotension In Obstetrics: Prevention And Therapy. *Acta Clinica Croatica*, 58(Suppl 1), 90–95. <https://doi.org/https://doi.org/10.20471/acc.2019.58.s1.13>
- Sugiyono. (2011). *Metode Penelitian Kuantitatif, Kualitatif dan R&D*. Bandung: Afabeta.
- Syahrum, & Salim. (2012). *Metodologi Penelitian Kuantitatif*. Bandung : Cipustaka Media.
- Tamamilang, C., Kandou, G. D., & Nelwan, J. E. (2018). Hubungan antara umur dan aktivitas fisik dengan derajat hipertensi di Kota Bitung Sulawesi Utara. *Jurnal KESMAS*, 7(5).
- Tanambel, P., Kumaat, L., & Lalenoh, D. (2017). Profil Penurunan Tekanan Darah (otensi) pada Pasien Sectio Caesarea yang Diberikan Anestesi Spinal dengan Menggunakan Bupivakain. *Jurnal E-Clinic*, 5(1) : 73.
- Taufik, M. (2017). Faktor Faktor Yang Mempengaruhi Kejadian Hipotensi Pada Anestesi Spinal RSUD KRT Setjonegoro Wonosobo. *Skripsi, Poltekkes*.
- Velde, M. Van De. (2019). Low-dose spinal anesthesia for cesarean section to prevent spinal-induced hypotension. *Co-Anesthesiology*, 32(3), 268–270. <https://doi.org/10.1097/ACO.0000000000000712>
- Wang, H. Z., Chen, H. W., Fan, Y. T., Jing, Y. L., Song, X. R., & She, Y. J. (2018). Relationship between body mass index and spread of spinal anesthesia in pregnant women: A randomized controlled trial. *Medical Science Monitor*, 24, 6144–6150. <https://doi.org/10.12659/MSM.909476>
- Whizar-Lugo, V. M. (2014). *Topics in Spinal Anaesthesia*. Institutos Nacionales de Salud.
- Widya, N. (2018). Hubungan Pemberian Preloading Cairan dengan Status Hemodinamik Pasien Anestesi Spinal di RSUD Muntilan. *Skripsi Thesis, Poltekkes Kemenkes Yogyakarta*.
- Zhafirah, A. (2019). *Pengaruh Indeks Massa Tubuh terhadap Kejadian Hipotensi Komplikasi Anestesi Spinal pada Pasien Sectio Caesarea di RSIA Siti Hawa Padang*. Dapat diakses di <http://scholar.unand.ac.id/43649/>.
- Zhu, Z., Tang, Y., Zhuang, J., Liu, Y., Wu, X., Cai, Y., ... Chen, P. (2019). Physical activity, screen viewing time, and overweight/obesity among Chinese children and adolescents: an update from the 2017 physical activity and fitness in China—the youth study. *BMC Public Health*, <https://doi.org/10.1186/s12889-019-6515-9>.